arXiv:1301.3766v1 [math.PR] 16 Jan 2013

Random directed forest and the Brownian web

Rahul Roy, Kumarjit Saha and Anish Sarkar [*]
Indian Statistical Institute, New Delhi

Abstract

Consider the d dimensional lattice $\mathbb{Z}^d$ where each vertex is *open* or *closed* with probability p or $1-p$ respectively. An open vertex $\mathbf{u} := (\mathbf{u}(1), \mathbf{u}(2), \ldots, \mathbf{u}(d))$ is connected by an edge to another open vertex which has the minimum L_1 distance among all the open vertices with $\mathbf{x}(d) > \mathbf{u}(d)$. It is shown that this random graph is a tree almost surely for $d = 2$ and 3 and it is an infinite collection of disjoint trees for $d \geq 4$. In addition for $d = 2$, we show that when properly scaled, family of its paths converges in distribution to the Brownian web.

Key words: Coexistence, Learning from neighbours, Markov chain, Random walk, Stationary distribution.

AMS 2000 Subject Classification: Primary: 60J10, 60K35; Secondary: 60C05, 62E10, 90B15, 91D30.

1 Introduction

Let $\Xi = \{\mathbf{x}_1, \mathbf{x}_2, \ldots\}$ be points of a Poisson point process on $\mathbb{R}^d$ of intensity 1 defined on some probability space $(\Omega, \mathcal{F}, P)$. For a point $\mathbf{x} = (\mathbf{x}(1), \ldots, \mathbf{x}(d)) \in \Xi$ let $h(\mathbf{x}) \in \Xi$ be such that (i) $\mathbf{x}(d) < h(\mathbf{x})(d)$ and (ii) $||\mathbf{x} - h(\mathbf{x})||_1 < ||\mathbf{x} - \mathbf{y}||_1$ for any $\mathbf{y} \in \Xi$ with $\mathbf{x}(d) < \mathbf{y}(d)$, where $|| \cdot ||_1$ denotes the L_1 metric on $\mathbb{R}^d$. This point $h(\mathbf{x})$ is almost surely unique. The directed spanning forest (DSF) is the random graph with vertex set Ξ and edge set $E := \{< \mathbf{x}, h(\mathbf{x}) >: \mathbf{x} \in \Xi\}$.

The study of DSF was initiated by Baccelli and Bordenave [BB07], Coupier and Tran [CT11] have shown that for $d = 2$ the DSF is a tree almost surely. Ferrari, Landim and Thorrisson [FLT04] also study a directed random graph on a Poisson point process, however the mechanism used to construct edges incorporate more

---

[*] E-Mail: `rahul, kumarjit10r, anish@isid.ac.in`

independence than is available in the DSF. They show that their random graph is a connected tree in dimensions 2 and 3, and a forest in dimensions 4 and more. In addition, for this random graph Ferrari, Fontes and Wu [FFW05] have showed that for $d = 2$ under a suitable diffusive scaling the random graph converges to a Brownian web.

In addition to DSF arising from a Poisson point process, there is also a construction of a similar process on the lattice. Let $\{U_{\mathbf{v}} : \mathbf{v} \in \mathbb{Z}^d\}$ be a collection of i.i.d. Uniform $(0, 1)$ random variables. Fix $0 < p < 1$ and let $V := \{\mathbf{v} \in \mathbb{Z}^d : U_{\mathbf{v}} < p\}$ be the set of *open* vertices of $\mathbb{Z}^d$. Given $\mathbf{u} \in \mathbb{Z}^d$, let $\mathbf{v} \in V$ be such that

1. $\mathbf{u}(d) < \mathbf{v}(d)$,
2. there does not exist any $\mathbf{w} \in V$ with $\mathbf{w}(d) > \mathbf{u}(d)$ such that $||\mathbf{u} - \mathbf{w}||_1 < ||\mathbf{u} - \mathbf{v}||_1$, and
3. for all $\mathbf{w} \in V$ with $\mathbf{w}(d) > \mathbf{u}(d)$ and $||\mathbf{u} - \mathbf{w}||_1 = ||\mathbf{u} - \mathbf{v}||_1$ we have $U_{\mathbf{v}} \leq U_{\mathbf{w}}$.

Such a $\mathbf{v}$ is almost surely unique and we denote it by $h(\mathbf{u})$. Let $< \mathbf{u}, h(\mathbf{u}) >$ be the edge joining $\mathbf{u}$ and $h(\mathbf{u})$ and let E denotes the edge set given by,

$$E := \{< \mathbf{u}, h(\mathbf{u}) >: \mathbf{u} \in V\}. \tag{1}$$



Figure 1: The construction of $h(\mathbf{u})$ from $\mathbf{u}$ on $\mathbb{Z}^2$. The shaded points are open, while the others are closed. Note that in order to get $h(\mathbf{u})$ from $\mathbf{u}$, we aquire information on the uniform values of the gray vertices.

The discrete DSF is the random graph $G := (V, E)$. These processes are called drainage processes (see Scheidegger [S67]) and have been studied in the physics literature extensively (Rodriguez-Iturbe [RR97]). Mathematically, for similar discrete processes but with a condition for constructing edges which allow more independence, the dichotomy in dimensions of having a single connected tree vis-a-vis a forest have been studied (see [GRS04], [ARS08]). In addition, their diffusive scaling limits have also been shown to be Brownian webs (see Coletti [CFD09]).

In this paper we study the most general discrete drainage network as defined in the preceding paragraph, and show that the tree/forest dichotomy in dimension holds, as well as the two dimensional scaling limit of the random graph is a Brownian web. Thus, this paper may be viewed as an extension, albeit in the discrete setting, of Coupier and Tran [CT11] result to any dimension. Our proof is different from that of Coupier and Tran; while their argument is percolation theoretic and crucially depends on the planarity of $\mathbb{R}^2$, our argument exploits a Markovian structure inherent in the DSF which allows us to extend the result to any dimension.

Theorem 1 *For $d = 2$ and $d = 3$ the random graph G is connected almost surely and consists of a single tree while for $d \geq 4$, it is a disconnected forest with each connected component being an infinite tree almost surely.*

Baccelli and Bordenave [BB07] have shown that scaled paths of the successive ancestors in the DSF converges weakly to the Brownian motion and also conjectured that the scaling limit of the DSF is a Brownian web. For the diffusive scaling of the discrete random graph G on $\mathbb{Z}^2$, let $\pi^{\mathbf{u}} := \{(\pi^{\mathbf{u}}(t), t) : t \geq \mathbf{u}(2)\}$, $\mathbf{u} \in \mathbb{Z}^d$ denote the graph of the path obtained by the edges $\{< h^{k-1}(\mathbf{u}), h^k(\mathbf{u}) >: k \geq 0)\}$ of the DSF $h^0(\mathbf{u}) := \mathbf{u}$. Let

$$\mathcal{X} := \{\pi^{\mathbf{u}} : \mathbf{u} \in V\},$$

and, for $\gamma, \sigma > 0$, define the n th order diffusive scaling of $\mathcal{X}$ by

$$\mathcal{X}_n(\gamma, \sigma) := \{(\frac{y(1)}{n\sigma}, \frac{y(2)}{n^2\gamma}) : \mathbf{y} := (y(1), y(2)) \in \mathcal{X}\}. \tag{2}$$

We have

Theorem 2 *For $d = 2$, there exist $\sigma := \sigma(p)$ and $\gamma := \gamma(p)$ such that as $n \to \infty$, $\mathcal{X}_n(\gamma, \sigma)$ converges weakly to the standard Brownian Web.*

The proof of Theorem 1 requires that we obtain a Markovian structure in our model and define suitable stop times for this Markov process. From these stop times the process regenerates which allows us to phrase the problem as a question of recurrence or transience of the Markov chain. This we do by obtaining a martingale for $d = 2$, using a Lyapunov function technique for $d = 3$ and a suitable coupling with a random walk with independent steps for $d = 4$.

The martingale obtained for $d = 2$ together with the fact that the distribution of the stop time has an exponentially decaying tail is used to prove Theorem 2.

The paper is structured as follows – in the next section we construct the paths of the graph G starting from two distinct vertices $\mathbf{u}$ and $\mathbf{v}$. In Section 3 we prove Theorem 1 and in Section 4, we prove Theorem 2.

2 Construction

We first detail a construction of the graph G which brings out a Markovian structure. Towards this end from two vertices $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^d$ we obtain the vertices $h^1(\mathbf{u}), h^2(\mathbf{u}), \dots$ and $h^1(\mathbf{v}), h^2(\mathbf{v}), \dots$ as a stochastic process as follows:-
Fix $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^d$ with $\mathbf{u}(d) = \mathbf{v}(d)$ and set $\tilde{H}_0(\mathbf{u}, \mathbf{v}) = \emptyset$. Let

$$\begin{aligned} & g_1(\mathbf{u}) := h(\mathbf{u}), g_1(\mathbf{v}) := h(\mathbf{v}) \text{ and} \\ & \tilde{H}_1(\mathbf{u}, \mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}) : \mathbf{w} \in \mathbb{Z}^d, \mathbf{w}(d) > \min\{g_1(\mathbf{u})(d), g_1(\mathbf{v})(d)\} \text{ and} \\ & \qquad\qquad ||\mathbf{u} - \mathbf{w}||_1 \le ||\mathbf{u} - g_1(\mathbf{u})||_1 \text{ or } ||\mathbf{v} - \mathbf{w}||_1 \le ||\mathbf{v} - g_1(\mathbf{v})||_1\} \end{aligned}$$



Figure 2: The construction of $g_1(\mathbf{u}), g_1(\mathbf{v})$. The union of the two shaded region is $\tilde{H}_1(\mathbf{u}, \mathbf{v})$.

Having obtained $g_n(\mathbf{u})$, $g_n(\mathbf{v})$ and the history set $\tilde{H}_n(\mathbf{u}, \mathbf{v})$, to obtain $g_{n+1}(\mathbf{u})$, $g_{n+1}(\mathbf{v})$ and $\tilde{H}_{n+1}(\mathbf{u}, \mathbf{v})$ we need to explain the evolution of the process for the three cases

$$(i)\ g_n(\mathbf{u})(d) = g_n(\mathbf{v})(d), \quad (ii)\ g_n(\mathbf{u})(d) > g_n(\mathbf{v})(d) \text{ and } (iii)\ g_n(\mathbf{u})(d) < g_n(\mathbf{v})(d).$$

In case (i), both the vertices $g_n(\mathbf{u})$ and $g_n(\mathbf{v})$ move, while in cases (ii) and (iii), the vertex with the smaller d th coordinate moves and the other vertex remains fixed, albeit with a new label. The history set $H_n(\mathbf{u}, \mathbf{v})$ is obtained accordingly. Formally, in case (i), i.e., $g_n(\mathbf{u})(d) = g_n(\mathbf{v})(d)$, set

$$\begin{aligned}
&g_{n+1}(\mathbf{u}) := h(g_n(\mathbf{u})), \quad g_{n+1}(\mathbf{v}) := h(g_n(\mathbf{v})) \text{ and}\\
&\tilde{H}_{n+1}(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}) : \mathbf{w} \in \mathbb{Z}^d, \mathbf{w}(d) > \min\{g_{n+1}(\mathbf{u})(d), g_{n+1}(\mathbf{v})(d)\} \text{ and}\\
&\qquad\qquad \text{at least one of the following three hold:}\\
&\qquad\qquad\qquad \text{(a) } ||g_n(\mathbf{u}) - \mathbf{w}||_1 \le ||g_n(\mathbf{u}) - g_{n+1}(\mathbf{u})||_1,\\
&\qquad\qquad\qquad \text{(b) } ||g_n(\mathbf{v}) - \mathbf{w}||_1 \le ||g_n(\mathbf{v}) - g_{n+1}(\mathbf{v})||_1,\\
&\qquad\qquad\qquad \text{(c) } \mathbf{w} \in \tilde{H}_n(\mathbf{u},\mathbf{v})\};
\end{aligned}$$



Figure 3: The construction of the process $\{g_j(\mathbf{u}), g_j(\mathbf{v}), \tilde{H}_j(\mathbf{u},\mathbf{v}) : j \ge 0\}$ from $\mathbf{u}$ and $\mathbf{v}$. Case (i): at n th step if $g_n(\mathbf{u})(d) = g_n(\mathbf{v})(d)$ then both of them move together.

in case (ii) i.e., $g_n(\mathbf{u})(d) > g_n(\mathbf{v})(d)$, set

$$\begin{aligned}
&g_{n+1}(\mathbf{u}) := g_n(\mathbf{u}), g_{n+1}(\mathbf{v}) := h(g_n(\mathbf{v})) \text{ and}\\
&\tilde{H}_{n+1}(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}) : \mathbf{w} \in \mathbb{Z}^d, \mathbf{w}(d) > \min\{g_{n+1}(\mathbf{u})(d), g_{n+1}(\mathbf{v})(d)\} \text{ and}\\
&\qquad\qquad \text{either } ||g_n(\mathbf{v}) - \mathbf{w}||_1 \le ||g_n(\mathbf{v}) - g_{n+1}(\mathbf{v})||_1 \text{ or } \mathbf{w} \in \tilde{H}_n(\mathbf{u},\mathbf{v})\};
\end{aligned}$$

in case (iii) i.e., $g_n(\mathbf{u})(d) < g_n(\mathbf{v})(d)$, set

$$
\begin{aligned}
& g_{n+1}(\mathbf{u}) := h(g_n(\mathbf{u})), g_{n+1}(\mathbf{v}) := g_n(\mathbf{v}) \text{ and} \\
& \tilde{H}_{n+1}(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}) : \mathbf{w} \in \mathbb{Z}^d, \mathbf{w}(d) > \min\{g_{n+1}(\mathbf{u})(d), g_{n+1}(\mathbf{v})(d)\} \text{ and} \\
& \qquad\qquad \text{either } ||g_n(\mathbf{u}) - \mathbf{w}||_1 \leq ||g_n(\mathbf{u}) - g_{n+1}(\mathbf{u})||_1 \text{ or } \mathbf{w} \in \tilde{H}_n(\mathbf{u},\mathbf{v})\}.
\end{aligned}
$$


$\tilde{H}_n(\mathbf{u},\mathbf{v})$
$g_n(\mathbf{u})$
$g_n(\mathbf{v})$
$\tilde{H}_{n+1}(\mathbf{u},\mathbf{v})$
$g_{n+1}(\mathbf{u}) = g_n(\mathbf{u})$
$g_{n+1}(\mathbf{v})$
$g_n(\mathbf{v})$

Figure 4: The construction of the process $\{g_j(\mathbf{u}), g_j(\mathbf{v}), \tilde{H}_j(\mathbf{u},\mathbf{v}) : j \geq 0\}$ from $\mathbf{u}$ and $\mathbf{v}$. Case (ii): at n th step if $g_n(\mathbf{u})(d) \neq g_n(\mathbf{v})(d)$ then vertex with smaller d coordinate moves.

Note that $\{g_n(\mathbf{u}), g_n(\mathbf{v}) : n \geq 0\}$ is a relabelling of the vertices $\{h^k(\mathbf{u}), h^k(\mathbf{v}) : k \geq 0\}$, the indexing is done to ensure that the processes from $\mathbf{u}$ and $\mathbf{v}$ move in tandem.

Given $g_n(\mathbf{u}), g_n(\mathbf{v})$ and $\tilde{H}_n(\mathbf{u},\mathbf{v})$, with $g_n(\mathbf{u})(d) < g_n(\mathbf{v})(d)$, to obtain $g_{n+1}(\mathbf{u})$ and $g_{n+1}(\mathbf{v})$ we need to know (i) $U_{\mathbf{w}}$ for vertices $\mathbf{w}$ on the boundary of the region $\{\mathbf{x} : (\mathbf{x}, U_{\mathbf{x}}) \in \tilde{H}_n(\mathbf{u},\mathbf{v})\}$, and (ii) $U_{\mathbf{w}}$ for vertices $\mathbf{w} \in \mathbb{Z}^d \setminus \tilde{H}_n(\mathbf{u},\mathbf{v})$ with $\mathbf{w}(d) > g_n(\mathbf{u})(d)$. This latter part we could check independently in a sequential manner by studying the unexplored vertices $\mathbf{x} \notin \tilde{H}_n(\mathbf{u},\mathbf{v})$ with $||\mathbf{x} - g_n(\mathbf{u})||_1 = k$, for $k = 1, 2, \ldots$, and stopping the first time we obtain a k with $U_{\mathbf{x}} < p$. A similar method may be employed when $g_n(\mathbf{u})(d) = g_n(\mathbf{v})(d)$, with the sequential procedure used for both $g_n(\mathbf{u})$ and $g_n(\mathbf{v})$. Thus $g_{n+1}(\mathbf{u}), g_{n+1}(\mathbf{v})$ and $\tilde{H}_{n+1}(\mathbf{u},\mathbf{v})$ depends only on $g_n(\mathbf{u}), g_n(\mathbf{v})$ and $\tilde{H}_n(\mathbf{u},\mathbf{v})$ together with the values of the uniform random variables associated with the unexplored vertices. This observation yields the Markov property:

Proposition 1 *For* $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^d$, $\mathbf{u}(d) = \mathbf{v}(d)$ *the process* $\{(g_n(\mathbf{u}), g_n(\mathbf{v}), \tilde{H}_n(\mathbf{u}, \mathbf{v})),$ $n \geq 0\}$ *is Markov with state space* $\mathbb{Z}^d \times \mathbb{Z}^d \times \{(\mathbf{w}, U_{\mathbf{w}}) : \mathbf{w} \in S,\ S \subset \mathbb{Z}^d,\ S \text{ finite}\}$.

Let $\{X_n(\mathbf{u}, \mathbf{v}), H_n(\mathbf{u}, \mathbf{v}) : n \geq 0\}$ be a time homogeneous modification of the above Markov process where, for $g_n(\mathbf{u})(d) > g_n(\mathbf{v})(d)$

$$\begin{aligned} X_n(\mathbf{u}, \mathbf{v}) &:= g_n(\mathbf{u}) - g_n(\mathbf{v}) \\ H_n(\mathbf{u}, \mathbf{v}) &:= \{(\mathbf{w} - g_n(\mathbf{v}), U_{\mathbf{w}}) : (\mathbf{w}, U_{\mathbf{w}}) \in \tilde{H}_n(\mathbf{u}, \mathbf{v})\}, \end{aligned}$$

and, for $g_n(\mathbf{u})(d) \leq g_n(\mathbf{v})(d)$

$$\begin{aligned} X_n(\mathbf{u}, \mathbf{v}) &:= g_n(\mathbf{v}) - g_n(\mathbf{u}) \\ H_n(\mathbf{u}, \mathbf{v}) &:= \{(\mathbf{w} - g_n(\mathbf{u}), U_{\mathbf{w}}) : (\mathbf{w}, U_{\mathbf{w}}) \in \tilde{H}_n(\mathbf{u}, \mathbf{v})\}. \end{aligned}$$

Remark: Although we have defined the Markov process starting from two distinct vertices, for any $k \geq 1$ a similar method can be used to define the Markov process $\{g_n(\mathbf{u}_1), \dots, g_n(\mathbf{u}_k), H_n(\mathbf{u}_1, \dots, \mathbf{u}_k) : n \geq 0\}$ starting from k distinct vertices $\mathbf{u}_1, \dots, \mathbf{u}_k$ with $\mathbf{u}_1(d) = \mathbf{u}_2(d) = \cdots = \mathbf{u}_k(d)$.

Let

$$L_n(\mathbf{u}, \mathbf{v}) := \begin{cases} 0 & \text{if } H_n(\mathbf{u}, \mathbf{v}) = \emptyset \\ \max\{\mathbf{x}(d) : (\mathbf{x}, y) \in H_n(\mathbf{u}, \mathbf{v}) \text{ for some } y \in [0, 1]\} & \text{otherwise} \end{cases}$$

denote the height of the history set $H_n(\mathbf{u}, \mathbf{v})$.

Before we proceed we note some properties of the history set. Given a vertex $\mathbf{w} \in \mathbb{Z}^d$, let $\mathbf{w}_m$ denote the vertex

$$\mathbf{w}_m(i) := \begin{cases} \mathbf{w}(i) & \text{for } 1 \leq i \leq d - 1, \\ m + \mathbf{w}(d) & \text{for } i = d \end{cases}$$

and suppose $g_n(\mathbf{u})(d) \leq g_n(\mathbf{v})(d)$.

(a) The history region $T_n(\mathbf{u}, \mathbf{v}) := \{\mathbf{w} : (\mathbf{w}, U_{\mathbf{w}}) \in \tilde{H}_n(\mathbf{u}, \mathbf{v})\}$ is a finite union of d-dimensional tetrahedrons, with each tetrahedron in this region having a $d - 1$ dimensional cube as a base on the hyperplane $\{\mathbf{x} \in \mathbb{Z}^d : \mathbf{x}(d) = 1 + g_n(\mathbf{u})(d)\}$. Also, $U_{\mathbf{w}} \geq p$ for every vertex $\mathbf{w}$ in the interior of the region $T_n(\mathbf{u}, \mathbf{v})$.

(b) The tallest tetrahedron in the history region $T_n(\mathbf{u}, \mathbf{v})$ is of height $L_n(\mathbf{u}, \mathbf{v})$.

(c) The vertices $\{(g_n(\mathbf{u}))_m : m \geq 1\}$ and $\{(g_n(\mathbf{v}))_m : m \geq 1\}$ are *not* in $T_n(\mathbf{u}, \mathbf{v})$. (This is represented for the history region $\tilde{H}_{n+1}(\mathbf{u}, \mathbf{v})$ in Figure 5.)



Figure 5: Geometry of the history region $T_{n+1}(\mathbf{u},\mathbf{v})$ in $\mathbb{Z}^2$: vertices $\{(g_{n+1}(\mathbf{u}))_m : m \geq 1\}$ and $\{(g_{n+1}(\mathbf{v}))_m : m \geq 1\}$ are not in $T_{n+1}(\mathbf{u},\mathbf{v})$.

Let

$$\tau(\mathbf{u},\mathbf{v}) := \inf\{n \geq 1 : H_n(\mathbf{u},\mathbf{v}) = \emptyset\}.$$

Note that $\tau(\mathbf{u},\mathbf{v}) = \inf\{n \geq 1 : L_n(\mathbf{u},\mathbf{v}) = 0\}$ denotes the number of steps required for the process to regenerate, i.e. to reach a state of empty history. Also, at the regeneration step, the two paths must be at the same level in terms of their d th coordinate, i.e. $g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{u})(d) = g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{v})(d)$. (See Figure 6.)


$g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{u})$
$g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{v})$
$\mathbf{u}$
$\mathbf{u}$

Figure 6: At the regeneration step $\tau(\mathbf{u},\mathbf{v})$ of the process $\{g_n(\mathbf{u}), g_n(\mathbf{v}), H_n(\mathbf{u},\mathbf{v}); n \geq 0\}$ we have $g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{u})(d) = g_{\tau(\mathbf{u},\mathbf{v})}(\mathbf{v})(d)$.

Proposition 2 *There exist constants $C_1, C_2 > 0$ such that, for all $n \geq 1$,*

$$\mathbb{P}(\tau(\mathbf{u},\mathbf{v}) \geq n | \mathbf{u}(d) = \mathbf{v}(d), H_0(\mathbf{u},\mathbf{v}) = \emptyset) \leq C_1 \exp(-C_2 n).$$

Proof: First note that property (c) of the history region ensures that, for every $n \geq 0$, given $\tilde{H}_n(\mathbf{u},\mathbf{v})$, the random variables

$$J_n(\mathbf{u}) := \inf\{m : g_n(\mathbf{u})_m \in V\} \text{ and } J_n(\mathbf{v}) := \inf\{m : g_n(\mathbf{v})_m \in V\} \tag{3}$$

are geometric random variables, each with success probability p.

Also, let $\{G_n : n \geq 0\}$ be a family of i.i.d. geometric random variables with success probability p, independent of $\{U_{\mathbf{x}} : \mathbf{x} \in \mathbb{Z}^d\}$.

We construct a sequence of random variables $J_0, J_1, \ldots$ where, for $n \geq 0$,

$$J_n = \begin{cases} \max\{J_n(\mathbf{u}), J_n(\mathbf{v})\} \text{ if } g_n(\mathbf{u})(d) = g_n(\mathbf{v})(d) \text{ and } g_n(\mathbf{u}) \neq g_n(\mathbf{v}), \\ \max\{J_n(\mathbf{u}), G_n\} \text{ if } g_n(\mathbf{u})(d) < g_n(\mathbf{v})(d) \text{ or } g_n(\mathbf{u}) = g_n(\mathbf{v}), \\ \max\{J_n(\mathbf{v}), G_n\} \text{ if } g_n(\mathbf{u})(d) > g_n(\mathbf{v})(d). \end{cases}$$

We have

$$\mathbb{P}(J_{n+1} \leq m | g_n(\mathbf{u}), g_n(\mathbf{v}), H_n(\mathbf{u},\mathbf{v})) = \mathbb{P}(\max\{G_1, G_2\} \leq m) \text{ for all } m. \tag{4}$$

Consider the Markov process $\{M_n(\mathbf{u},\mathbf{v}), g_n(\mathbf{u}), g_n(\mathbf{u}), \tilde{H}_n(\mathbf{u},\mathbf{v}) : n \geq 0\}$ where $g_n(\mathbf{u}), g_n(\mathbf{u}), \tilde{H}_n(\mathbf{u},\mathbf{v})$ are defined as earlier and $M_n(\mathbf{u},\mathbf{v})$ is a non-negative integer valued random variable defined recursively as

$$\begin{aligned} M_0(\mathbf{u},\mathbf{v}) &:= 0 \text{ and} \\ M_n(\mathbf{u},\mathbf{v}) &:= \begin{cases} \max\{M_{n-1}(\mathbf{u},\mathbf{v}) - 1, 0\} & \text{if } J_{n-1} = 1, \\ M_{n-1}(\mathbf{u},\mathbf{v}) & \text{if } 1 < J_{n-1} \leq M_{n-1}(\mathbf{u},\mathbf{v}) \\ & \text{and } M_{n-1}(\mathbf{u},\mathbf{v}) \geq l_0, \\ J_{n-1} & \text{if } J_{n-1} > M_{n-1}(\mathbf{u},\mathbf{v}) \geq l_0, \\ l_0 + J_{n-1} & \text{otherwise;} \end{cases} \end{aligned} \tag{5}$$

here $l_0 \geq 1$ is a constant which is specified later. First note that $L_0(\mathbf{u},\mathbf{v}) = M_0(\mathbf{u},\mathbf{v}) = 0$ and, by induction, we may show that the coupling above ensures $0 \leq L_n(\mathbf{u},\mathbf{v}) \leq M_n(\mathbf{u},\mathbf{v})$ for all $\mathbf{u},\mathbf{v}$ and all $n \geq 0$. Indeed, assuming $L_n(\mathbf{u},\mathbf{v}) \leq M_n(\mathbf{u},\mathbf{v})$, we have

(i) if $J_n = 1$ then $L_{n+1}(\mathbf{u},\mathbf{v}) = \max\{L_n(\mathbf{u},\mathbf{v}) - 1, 0\} \leq \max\{M_n(\mathbf{u},\mathbf{v}) - 1, 0\} = M_{n+1}(\mathbf{u},\mathbf{v})$;

(ii) if $M_n(\mathbf{u},\mathbf{v}) \geq l_0$ and $1 < J_n \leq M_n(\mathbf{u},\mathbf{v})$, then $M_{n+1}(\mathbf{u},\mathbf{v}) = M_n(\mathbf{u},\mathbf{v})$, while $L_{n+1}(\mathbf{u},\mathbf{v}) \leq \max\{J_n, L_n(\mathbf{u},\mathbf{v})\} \leq M_{n+1}(\mathbf{u},\mathbf{v})$;

(iii) if $J_n > M_n(\mathbf{u},\mathbf{v}) \geq l_0$, then $J_n > L_n(\mathbf{u},\mathbf{v})$ and so $L_{n+1}(\mathbf{u},\mathbf{v}) \leq J_n = M_{n+1}(\mathbf{u},\mathbf{v})$;

(iv) if $L_n(\mathbf{u},\mathbf{v}) \leq M_n(\mathbf{u},\mathbf{v}) < l_0$ then
$L_{n+1}(\mathbf{u},\mathbf{v}) \leq L_n(\mathbf{u},\mathbf{v}) + J_n \leq l_0 + J_n = M_{n+1}(\mathbf{u},\mathbf{v})$.

Hence, to prove the proposition, it suffices to show that, taking $\tau(M(\mathbf{u},\mathbf{v})) := \inf\{n : M_n(\mathbf{u},\mathbf{v}) = 0\}$, we have

$$\mathbb{P}(\tau(M(\mathbf{u},\mathbf{v})) \geq n | M_0(\mathbf{u},\mathbf{v}) = 0) \leq C_1 \exp(-C_2 n), \tag{6}$$

for some constants $C_1, C_2 > 0$.

Towards this end we consider a random walk $S_n := S_0 + \sum_{i=1}^n Z_i$, where Z_i, $i \geq 1$, are i.i.d. copies of the random variable $Z^{(l_0)}$ which takes values in the set $\{-1, 0, 1, 2, \ldots\}$ and is defined by

$$\mathbb{P}(Z^{(l_0)} = k) = \begin{cases} 2p(1-p)^{l_0+k} & \text{if } k = 1, 2, \ldots, \\ p^2 & \text{if } k = -1, \\ 1 - p^2 - 2(1-p)^{l_0+1} & \text{if } k = 0. \end{cases} \tag{7}$$

The random walk $\{S_n,\ n \geq 0\}$ has negative drift for l_0 such that

$$\sum_{k=1}^{\infty} 2kp(1-p)^{l_0+k} = 2\frac{(1-p)^{l_0+1}}{p} < p^2. \tag{8}$$

A comparison of the probabilities given in (5) and (7) shows that for $l \geq l_0$, $k \geq -1$ and for all $\mathbf{u}, \mathbf{v}$ with $\mathbf{u}(d) = \mathbf{v}(d)$

$$\mathbb{P}(M_{n+1}(\mathbf{u},\mathbf{v}) \geq l + k | M_n(\mathbf{u},\mathbf{v}) = l) \quad \leq \quad \mathbb{P}(S_{n+1} \geq l + k | S_n = l). \tag{9}$$

Thus the process $\{S_n,\ n \geq 0\}$ dominates the process $\{M_n(\mathbf{u},\mathbf{v}),\ n \geq 0\}$ in the sense that for any $\mathbf{u}, \mathbf{v}$ and for any $l \geq l_0$,

$$\mathbb{P}(\tau^S \geq n | S_0 = l) \geq \mathbb{P}(\tau_M^{l_0}(\mathbf{u},\mathbf{v}) \geq n | M_0(\mathbf{u},\mathbf{v}) = l); \tag{10}$$

where $\tau^S := \inf\{n \geq 1 : S_n = l_0 - 1\}$ and $\tau_M^{l_0}(\mathbf{u},\mathbf{v}) := \inf\{n : M_n(\mathbf{u},\mathbf{v}) = l_0 - 1\}$. By construction S_n is a left continuous random walk with negative drift and hence we know that

$$\mathbb{P}(\tau^S \geq n | S_0 = l_0 + \max\{G_1, G_2\}) \leq c_3 \exp(-c_4 n) \text{ for some constants } c_3, c_4 > 0.$$

Let $\Gamma_1, \Gamma_2, \ldots$ be i.i.d. random variables with

$$\mathbb{P}(\Gamma_1 \leq n) := P(\tau^S \leq n | S_0 = l_0 + \max\{G_1, G_2\}).$$

Suppose $G^{(l_0)}$ is a geometric random variable with success probability $p^{2(l_0-1)}$ and independent of $\Gamma_1, \Gamma_2, \ldots$. For

$$\Gamma_{rw} := \sum_{i=1}^{J^{(l_0)}-1} (\Gamma_i + l_0) + l_0,$$

we have

$$\mathbb{P}(\Gamma_{rw} \geq n) \leq c_5 \exp\left(-c_6 n\right) \text{ for some constants } c_5, c_6 > 0.$$

This completes the proof since from the Markov property it follows that Γ_{rw} stochastically dominates $\tau(X(\mathbf{u}, \mathbf{v}))$. $\square$

Before we end this section we make same observations of our construction which are used later.

1. For any two pairs of open vertices $\mathbf{u}_1, \mathbf{v}_1$ with $\mathbf{u}_1(d) = \mathbf{v}_1(d)$ and $\mathbf{u}_2, \mathbf{v}_2$ with $\mathbf{u}_2(d) = \mathbf{v}_2(d)$, $\tau(M(\mathbf{u}_1, \mathbf{v}_1))$ and $\tau(M(\mathbf{u}_2, \mathbf{v}_2))$ have the same distribution.

2. Suppose $\{G^1_i : i \geq 0\}$ and $\{G^2_i : i \geq 0\}$ are two independent families of i.i.d. geometric random variables each with success probability p and independent of $\{M_n(\mathbf{u}, \mathbf{v}) : n \geq 0\}$. For

   $$\Gamma := \sum_{i=0}^{\tau(M(\mathbf{u},\mathbf{v}))} (G^1_i + G^2_i),$$

   we have that

   $$P(\Gamma \geq n) \leq D_1 \exp -(D_2 n) \tag{11}$$

   for some constants $D_1, D_2 > 0$.

3. For $j \geq 0$ and $\mathbf{u}, \mathbf{v} \in V$ with $\mathbf{u}(d) = \mathbf{v}(d)$, let

   $$W_j(\mathbf{u}, \mathbf{v}) := \sum_{i=\tau_j(\mathbf{u},\mathbf{v})+1}^{\tau_{j+1}(\mathbf{u},\mathbf{v})} \{||g_i(\mathbf{u}) - g_{i-1}(\mathbf{u})||_1 + ||g_i(\mathbf{v}) - g_{i-1}(\mathbf{v})||_1\} \tag{12}$$

   denote the width of the region explored between the regeneration times $\tau_j(\mathbf{u}, \mathbf{v})$ and $\tau_{j+1}(\mathbf{u}, \mathbf{v})$.

   Similarly, for the process starting from only one vertex $\mathbf{u}$, the width of the region explored between the j th regeneration step and $j+1$ th regeneration step is

   $$W_j(\mathbf{u}) := \sum_{i=\tau_j(\mathbf{u})+1}^{\tau_{j+1}(\mathbf{u})} \{||g_i(\mathbf{u}) - g_{i-1}(\mathbf{u})||_1\}, \tag{13}$$

where $\tau_j(\mathbf{u})$ is the j th regeneration step for the process $\{h^n(\mathbf{u}) : n \geq 0\}$.

From property (c) of the history region we have that, for any $a > 0$,

$$\begin{aligned} \mathbb{P}(W_j(\mathbf{u},\mathbf{v}) \geq a) &\leq \mathbb{P}(\Gamma \geq a) \leq D_1 \exp -(D_2 a), \qquad (14)\\ \mathbb{P}(W_j(\mathbf{u}) \geq a) &\leq \mathbb{P}(\Gamma \geq a) \leq D_1 \exp -(D_2 a). \qquad (15) \end{aligned}$$

3 Tree and Forest

In this section we prove Theorem 1. For $\mathbf{x} \in \mathbb{Z}^d$ let $\overline{\mathbf{x}} := (\mathbf{x}(1), \ldots, \mathbf{x}(d-1)) \in \mathbb{Z}^{d-1}$. For any $\mathbf{u}, \mathbf{v} \in V$, with $\mathbf{u}(d) = \mathbf{v}(d)$,

$$\{Z_j(\mathbf{u},\mathbf{v}) := \overline{g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v})} - \overline{g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v})} : j \geq 0\} \qquad (16)$$

is a time homogeneous Markov chain with a single absorbing state at $\bar{\mathbf{0}}$ on $\mathbb{Z}^{d-1}$.

3.1 d=2

To begin with, for $d = 2$, we first obtain a martingale which we also use later for the proof of Theorem 2. Let $\mathbf{u}, \mathbf{v} \in V$ with $\mathbf{u}(2) = \mathbf{v}(2)$ and consider the filtration $\mathcal{F}_0 \subseteq \mathcal{F}_1 \subseteq \cdots$ where $\mathcal{F}_i := \sigma\{U_{\mathbf{x}} : \mathbf{u}(2) \leq \mathbf{x}(2) \leq \mathbf{u}(2) + i\}$. Let $T_j(\mathbf{u},\mathbf{v}) := g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(d) - \mathbf{u}(d)$ be the distance travelled in the d-th direction till the j th regeneration time. Note that in our earlier notation $\tau_j(\mathbf{u},\mathbf{v})$ denotes the number of steps taken till the j-th regeneration. Thinking of the d-th direction as time, $T_j(\mathbf{u},\mathbf{v})$ denotes the time taken for j-th regeneration and it is easy to see that $T_j(\mathbf{u},\mathbf{v})$ is a stop time with respect to the filtration $\{\mathcal{F}_i : i \geq 0\}$.

Proposition 3 *For $d = 2$ and $\mathbf{u}, \mathbf{v} \in V$ with $\mathbf{u}(2) = \mathbf{v}(2)$, the process $\{g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) : j \geq 0\}$ is a martingale with respect to the filtration $\{\mathcal{F}_{T_j(\mathbf{u},\mathbf{v})} : j \geq 0\}$.*

Proof: Since $g_{\tau_{j+1}(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1)$ is $\mathcal{F}_{T_{j+1}(\mathbf{u},\mathbf{v})}$ measurable and independent of $\mathcal{F}_{T_j(\mathbf{u},\mathbf{v})}$, it suffices to show that $\mathbb{E}[g_{\tau_{j+1}(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1)] = 0$.

It is easy to see that the 1-dimensional process $\{h^{\tau_i(\mathbf{u})}(\mathbf{u})(1) : i \geq 0\}$ is a symmetric random walk with increments being independent and identically distributed. For $j \geq 0$ let $S_j(\mathbf{u})$ be the time taken for j-th regeneration of the process $\{h^n(\mathbf{u}) : n \geq 0\}$, i.e.

$$S_j(\mathbf{u}) := h^{\tau_j(\mathbf{u})}(\mathbf{u})(2) - \mathbf{u}(2).$$

Note, for any $\mathbf{u}, \mathbf{v}$, a regeneration time of the process $\{g_n(\mathbf{u}), g_n(\mathbf{v}), H_n(\mathbf{u},\mathbf{v}); n \geq 0\}$ must necessarily be a regeneration time of the process $\{h^n(\mathbf{u}) : n \geq 0\}$. Hence we have a random sequence $I_0 < I_1 < \cdots$ such that $S_{I_j}(\mathbf{u}) = T_j(\mathbf{u},\mathbf{v})$. Also for


$S_3(\mathbf{u})$
$S_2(\mathbf{u})$
$S_1(\mathbf{u})$
$h^{\tau_3(\mathbf{u})}(\mathbf{u})$
$h^{\tau_2(\mathbf{u})}(\mathbf{u})$
$h^{\tau_1(\mathbf{u})}(\mathbf{u})$
$T_1(\mathbf{u},\mathbf{v})$
$h^{\tau_2(\mathbf{v})}(\mathbf{v})$
$h^{\tau_1(\mathbf{v})}(\mathbf{v})$
$S_2(\mathbf{v})$
$S_1(\mathbf{v})$
$\mathbf{u}$
$\mathbf{v}$

Figure 7: A regeneration time of the process $\{g_n(\mathbf{u}), g_n(\mathbf{v}), H_n(\mathbf{u},\mathbf{v}); n \geq 0\}$ must necessarily be a regeneration time of the process $\{h^n(\mathbf{u}); n \geq 0\}$ as well as of the process $\{h^n(\mathbf{v}); n \geq 0\}$.

any $n \geq 1$ and $k \geq 1$

$$\begin{aligned}\{I_j \leq n\} \cap \{S_n(\mathbf{u}) \leq k\} \in \sigma\{U_{\mathbf{x}} : \mathbf{x} \in [\mathbf{u}(1) - k^2, \mathbf{u}(1) + k^2] \times [\mathbf{u}(2), \mathbf{u}(2) + k] \\ \cup [\mathbf{v}(1) - k^2, \mathbf{v}(1) + k^2] \times [\mathbf{v}(2), \mathbf{v}(2) + k]\} \subseteq \mathcal{F}_k\end{aligned}$$

Hence I_j is a stop time with respect to the filtration $\{\mathcal{F}_{S_k(\mathbf{u})} : k \geq 0\}$ and moreover from (11) it has moments of all orders. It is easy to see that $h^{\tau_{k+1}(\mathbf{u})}(\mathbf{u}) - h^{\tau_k(\mathbf{u})}(\mathbf{u})$ is $\mathcal{F}_{S_{k+1}(\mathbf{u})}$ measurable and independent of $\mathcal{F}_{S_k(\mathbf{u})}$. Thus using Wald's identity we have

$$\mathbb{E}[g_{\tau_{j+1}(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1)] = \mathbb{E}\left(\sum_{k=1}^{I_{j+1}-I_j} [h^{\tau_{I_j+k}(\mathbf{u})}(\mathbf{u})(1) - h^{\tau_{I_j+k-1}(\mathbf{u})}(\mathbf{u})(1)]\right) = 0.$$

$\square$

Now we will show that the paths originating from $\mathbf{u}$ and $\mathbf{v}$ meet almost surely. First suppose that $\mathbf{u}(2) = \mathbf{v}(2)$. Since the paths $\{g_n(\mathbf{u}) : n \geq 0\}$ and $\{g_n(\mathbf{v}) : n \geq 0\}$ do not cross each other $\{g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) : j \geq 0\}$ forms a nonnegative or nonpositive martingale depending on whether $\mathbf{v}(1) > \mathbf{u}(1)$ or not, and by the martingale convergence theorem, there exists a random variable Z_∞ such that $g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) \to Z_\infty$ a.s. as $j \to \infty$. Also, 0 being the only absorbing state of the Markov chain $\{g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v})(1) - g_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u})(1) : j \geq 0\}$ we have $Z_\infty = 0$ a.s., i.e., $g_n(\mathbf{u}) = g_n(\mathbf{v})$ a.s. for some $n \geq 0$.

Next suppose that $\mathbf{v}(2) > \mathbf{u}(2)$. Consider the random set

$$A = \{\mathbf{x} : \mathbf{x}(2) = \mathbf{u}(2), \mathbf{x}(1) \geq \mathbf{u}(1), \mathbf{x} \in V, h(\mathbf{x})(2) = \mathbf{v}(2), h(\mathbf{x})(1) = \mathbf{x}(1)\},$$

i.e. the set of vertices $\mathbf{x}$ at the same level as $\mathbf{u}$, but to the left of $\mathbf{u}$, for which $h(\mathbf{x})$ is at the same level as $\mathbf{v}$ and does not carry any history. For $n \geq 0$, E_n denotes the event that $\{(\mathbf{u}(1) + n, \mathbf{u}(2)) \in A$ and $(\mathbf{u}(1) + j, \mathbf{u}(2)) \notin A$ for all $0 \leq j < n\}$.

$$\begin{aligned}
&\mathbb{P}(g_n(\mathbf{u}) = g_m(\mathbf{v}) \text{ for some } m, n \geq 1) \\
&\geq \mathbb{P}(\text{ there exists } \mathbf{x} \in A,\ g_n(\mathbf{x}) = g_n(\mathbf{u}) \text{ for some } n \geq 0 \\
&\qquad\qquad\qquad\qquad \text{and } g_m(h(\mathbf{x})) = g_m(\mathbf{v}) \text{ for some } m \geq 0) \\
&= \sum_{n \geq 0} \mathbb{P}(g_n(\mathbf{x}) = g_n(\mathbf{u}) \text{ for some } n \geq 0,\ g_m(h(\mathbf{x})) = g_m(\mathbf{v}) \text{ for some } m \geq 0 | E_n)\mathbb{P}(E_n) \\
&= \sum_{n \geq 0} \mathbb{P}(E_n) \\
&= 1.
\end{aligned}$$

This completes the proof of Theorem 1 for $d = 2$.

3.2 $d = 3$

It suffices to show that the paths originating from $\mathbf{0}$ and $\mathbf{v}$ with $\mathbf{v}(3) = 0$ meet almost surely. For $d = 3$ we modify the $\mathbb{Z}^2$ valued Markov process $\{Z_j(\mathbf{0}, \mathbf{v}) : j \geq 0\}$ defined in (16) such that the transition probabilities are the same except that instead of $(0, 0)$ being an absorbing state we have

$$\mathbb{P}((0, 0) \to (0, 1)) = 1.$$

With a little abuse of notation we denote this modified chain also by $\{Z_j(\mathbf{0}, \mathbf{v}) : j \geq 0\}$. This modified chain is clearly irreducible. To show that the original Markov process is absorbed at $(0, 0)$, it suffices to show that the modified chain is recurrent. Consider the function $f : \mathbb{R}^2 \to [0, \infty)$ defined by

$$f(x) = \sqrt{\log(1 + ||x||_2^2)}.$$

Note that $f(x) \to \infty$ as $||x||_2 \to \infty$. To show that $\{Z_j(\mathbf{0}, \mathbf{v}) : j \geq 0\}$ is recurrent, by Foster's criterion [see Asmussen (1987), Proposition 5.3 of Chapter 1, page 18] it is enough to show that there exists $0 < k_0 < \infty$ such that

$$\mathbb{E}[f(Z_{j+1}(\mathbf{0}, \mathbf{v})) - f(Z_j(\mathbf{0}, \mathbf{v})) | Z_j(\mathbf{0}, \mathbf{v}) = \mathbf{x}] < 0 \text{ for all } \mathbf{x} \text{ with } ||\mathbf{x}||_2^2 > k_0. \quad (17)$$

Let $A_{\mathbf{x},k} := \{W_0(\mathbf{0},\mathbf{x}) \leq \frac{k}{4}\}$, where $W_0(\mathbf{0},\mathbf{x})$ is as in (12). The translation invariance of our model and the Markov property ensures that

$$
\begin{aligned}
&\mathbb{E}[f(Z_{j+1}(\mathbf{0},\mathbf{v})) - f(Z_j(\mathbf{0},\mathbf{v})) | Z_j(\mathbf{0},\mathbf{v}) = \mathbf{x}] \\
&= \mathbb{E}[f(Z_1(\mathbf{0},\mathbf{x})) - f(Z_0(\mathbf{0},\mathbf{x}))] \\
&= \mathbb{E}[f(Z_1(\mathbf{0},\mathbf{x})) - f(\mathbf{x})1_{A_{\mathbf{x},k}}] + \mathbb{E}[f(Z_1(\mathbf{0},\mathbf{x})) - f(\mathbf{x})1_{A^c_{\mathbf{x},k}}].
\end{aligned}
$$

Let $B_n(\mathbf{u}) := \{\mathbf{y} : ||\mathbf{y} - \mathbf{u}||_1 \leq n\}$. Now, for $||\mathbf{x}||_2 \geq k$, on the event $A_{\mathbf{x},k}$ the two boxes $D_0^{\mathbf{0},\mathbf{x}}(\mathbf{0}) := B_{W_0(\mathbf{0},\mathbf{x})}(\bar{\mathbf{0}}) \times [0, T_1(\mathbf{0},\mathbf{x})]$ and $D_0^{\mathbf{0},\mathbf{x}}(\mathbf{x}) := B_{W_0(\mathbf{0},\mathbf{x})}(\bar{\mathbf{x}}) \times [0, T_1(\mathbf{0},\mathbf{x})]$ are disjoint and contain the set of vertices explored till the first regeneration of the process $\{g_j(\mathbf{0}), g_j(\mathbf{x}), H_j(\mathbf{0},\mathbf{x}) : j \geq 0\}$. Moreover on this event, each of the two coordinates of the increments of the Markov process $\{Z_j(\mathbf{0},\mathbf{x}) : j \geq 0\}$ are symmetric about 0 and identically distributed. This observation together with the translation invariance property gives,


$g_{\tau_1(\mathbf{0},\mathbf{x})}(\mathbf{0})$
$T_1(\mathbf{0},\mathbf{x})$
$D_0^{\mathbf{0},\mathbf{x}}(\mathbf{0})$
$\mathbf{0}$
$g_{\tau_1(\mathbf{0},\mathbf{x})}(\mathbf{x})$
$D_0^{\mathbf{0},\mathbf{x}}(\mathbf{x})$
$\mathbf{x}$
$W_0(\mathbf{0},\mathbf{x})$

Figure 8: The shaded regions represent $D_0^{\mathbf{0},\mathbf{x}}(\mathbf{0})$ and $D_0^{\mathbf{0},\mathbf{x}}(\mathbf{x})$.

$$
\begin{aligned}
&m_r(k) := \mathbb{E}[(Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(1)1_{A_{\mathbf{x},k}}]^r = \mathbb{E}[(Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(2)1_{A_{\mathbf{x},k}}]^r \\
&m_{r,s}(k) := \mathbb{E}[((Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(1))^r((Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(2))^s 1_{A_{\mathbf{x},k}}] \\
&\qquad = \mathbb{E}[((Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(1))^r((Z_1(\mathbf{0},\mathbf{x}) - Z_0(\mathbf{0},\mathbf{x}))(2))^l 1_{A_{\mathbf{x},k}}] \\
&m_r(k) = 0 \text{ for all odd } r \\
&m_{r,s}(k) = 0 \text{ whenever either } r \text{ or } s \text{ is odd .}
\end{aligned}
$$

For $m \in \mathbb{N}$ on the event $A^c_{\mathbf{x},k} \cap \{W_1(\mathbf{0},\mathbf{x}) = \frac{k}{4} + m\}$ we have

$$
\begin{aligned}
||Z_1(\mathbf{0},\mathbf{x})||_2^2 - ||Z_0(\mathbf{0},\mathbf{x})||_2^2 &\leq ||g_{\tau_1(\mathbf{0},\mathbf{x})}(\mathbf{0})(1,2)||_1^2 + ||g_{\tau_1(\mathbf{0},\mathbf{x})}(\mathbf{v})(1,2) - \mathbf{v}(1,2)||_1^2 \\
&\leq 2(\frac{k}{4} + m)^2;
\end{aligned}
$$

hence, using (11),

$$\begin{aligned}\mathbb{E}[(||Z_1(\mathbf{0},\mathbf{x})||_2^2 - ||Z_0(\mathbf{0},\mathbf{x})||_2^2)1_{A_{\mathbf{x}}^c)}]^l &\leq \sum_{m\geq 1} 2(\frac{k}{4}+m)^{2l}\mathbb{P}(W_1(\mathbf{0},\mathbf{x}) = \frac{k}{4}+m)\\ &\leq \sum_{m\geq 1} 2(\frac{k}{4}+m)^{2l}\mathbb{P}(\Gamma = \frac{k}{4}+m)\\ &= o(k^{-2}) \text{ as } k\to\infty.\end{aligned}$$

The second inequality follows from (14). Now calculations similar to those in Theorem 2.1 in [GRS04] show that for $||\mathbf{x}||_2 \geq k$, as $k \to \infty$

$$\mathbb{E}[||Z_{j+1}(\mathbf{0},\mathbf{v})||_2^2 - ||Z_j(\mathbf{0},\mathbf{v})||_2^2 | Z_j(\mathbf{0},\mathbf{v}) = \mathbf{x}] = \alpha(k) + o(k^{-2}) \qquad (18)$$

$$\mathbb{E}[(||Z_{j+1}(\mathbf{0},\mathbf{v})||_2^2 - ||Z_j(\mathbf{0},\mathbf{v})||_2^2)^2 | Z_j(\mathbf{0},\mathbf{v}) = \mathbf{x}] \geq 2\alpha(k)||\mathbf{x}||_2^2 \qquad (19)$$

$$\mathbb{E}[(||Z_{j+1}(\mathbf{0},\mathbf{v})||_2^2 - ||Z_j(\mathbf{0},\mathbf{v})||_2^2)^3 | Z_j(\mathbf{0},\mathbf{v}) = \mathbf{x}] = O(||\mathbf{x}||_2^2), \qquad (20)$$

where $\alpha(k) := 2m_2(k)$. Using the above estimates in the Taylor series expansion of f, as in [GRS04], we get that there exists k_0 large enough so that for $||\mathbf{x}||_2 \geq k_0$

$$\mathbb{E}[f(Z_{j+1}(\mathbf{0},\mathbf{v})) - f(Z_j(\mathbf{0},\mathbf{v})) | Z_j(\mathbf{0},\mathbf{v}) = \mathbf{x}] < 0.$$

This proves that the modified Markov chain is recurrent and completes the proof.

3.3 $d \geq 4$

In this subsection we present the proof only for $d = 4$. The argument for $d > 4$ is similar. We first show that for $\mathbb{Z}^4$, $\mathbb{P}(G$ is disconnected $) > 0$. We start with two open vertices $\mathbf{u}$ and $\mathbf{v}$ in $\mathbb{Z}^4$ with $\mathbf{u}(4) = \mathbf{v}(4)$. Consider the $\mathbb{Z}^3$ valued Markov chain $\{Z_j(\mathbf{u},\mathbf{v}); j \geq 0\}$ where $Z_j(\mathbf{u},\mathbf{v})$ as in (16). We couple this Markov chain with a suitable random walk on $\mathbb{Z}^3$, and by comparing this chain with the random walk we show that there is a positive probability that the Markov chain does not get absorbed at $(0,0,0)$. We first construct the desired random walk and present the formal details of the coupling procedure thereafter.

Let $\{U_{\mathbf{x}}^a : \mathbf{x} \in \mathbb{Z}^4\}$ be a collection of i.i.d. $U(0,1)$ random variables and consider the directed spanning forest $G_a := (V_a, E_a)$ with the open vertices V_a and edges $E_a = \{< \mathbf{u}, h^a(\mathbf{u}) >: \mathbf{u} \in V_a\}$ obtained from $\{U_{\mathbf{x}}^a : \mathbf{x} \in \mathbb{Z}^4\}$, with $h^a(\mathbf{u})$ being similar to that in (1). Let G_b be another directed spanning forest obtained in a similar fashion from a collection $\{U_{\mathbf{x}}^b : \mathbf{x} \in \mathbb{Z}^4\}$ of i.i.d. $U(0,1)$ random variables independent of $\{U_{\mathbf{x}}^a : \mathbf{x} \in \mathbb{Z}^4\}$. Let $\{h_n^a(\mathbf{u}) : n \geq 0\}$ and $\{h_n^b(\mathbf{v}) : n \geq 0\}$ be the paths from $\mathbf{u}$ and $\mathbf{v}$ in the forests G_a and G_b respectively. Define

$$g_1^a(\mathbf{u}) := h_1^a(\mathbf{u}) \text{ and } g_1^b(\mathbf{v}) := h_1^b(\mathbf{v})$$

and

$$g_n^a(\mathbf{u}) := \begin{cases} h^a(g_{n-1}^a(\mathbf{u})) & \text{if } g_{n-1}^a(\mathbf{u})(4) \leq g_{n-1}^b(\mathbf{v})(4) \\ g_{n-1}^a(\mathbf{u}) & \text{otherwise,} \end{cases}$$

$$g_n^b(\mathbf{v}) := \begin{cases} h^b(g_{n-1}^b(\mathbf{v})) & \text{if } g_{n-1}^b(\mathbf{v})(4) \leq g_{n-1}^a(\mathbf{u})(4) \\ g_{n-1}^b(\mathbf{v}) & \text{otherwise.} \end{cases}$$

As in Section 2, this relabeling is done to ensure that the two processes move in tandem. Let $H_n^a(\mathbf{u},\mathbf{v})$ and $H_n^b(\mathbf{u},\mathbf{v})$ be the history sets associated with the process $\{g_n^a(\mathbf{u}), g_n^b(\mathbf{v}) : n \geq 0\}$, i.e.,

$$\begin{aligned} H_n^a(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}^a) : \mathbf{w} \in \mathbb{Z}^4, \mathbf{w}(4) > \min\{g_n^a(\mathbf{u})(4), g_n^b(\mathbf{v})(4)\} \text{ and} \\ \text{at least one of the following two hold:} \\ \text{(a)}||g_{n-1}^a(\mathbf{u}) - \mathbf{w}||_1 \leq ||g_{n-1}^a(\mathbf{u}) - g_n^a(\mathbf{u})||_1 \\ \text{(b)}\mathbf{w} \in H_{n-1}^a(\mathbf{u},\mathbf{v})\}; \\ H_n^b(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}^b) : \mathbf{w} \in \mathbb{Z}^4, \mathbf{w}(4) > \min\{g_n^a(\mathbf{u})(4), g_n^b(\mathbf{v})(4)\} \text{ and} \\ \text{at least one of the following two hold:} \\ \text{(a)}||g_{n-1}^b(\mathbf{v}) - \mathbf{w}||_1 \leq ||g_{n-1}^b(\mathbf{v}) - g_n^b(\mathbf{v})||_1 \\ \text{(b)}\mathbf{w} \in H_{n-1}^b(\mathbf{u},\mathbf{v})\}. \end{aligned}$$

Let $\tau^{ind}(\mathbf{u},\mathbf{v}) := \inf\{n : H_n^a(\mathbf{u},\mathbf{v}) = H_n^b(\mathbf{u},\mathbf{v}) = \emptyset\}$ denote the number of steps required for the regeneration of the process $\{g_n^a(\mathbf{u}), g_n^b(\mathbf{v}) : n \geq 0\}$. As in Proposition 2 we have for $\tau^{ind}(\mathbf{u},\mathbf{v})$

$$\mathbb{P}(\tau^{ind}(\mathbf{u},\mathbf{v}) \geq n | \mathbf{u}(4) = \mathbf{v}(4), H_0^a(\mathbf{u},\mathbf{v}) = H_0^b(\mathbf{u},\mathbf{v}) = \emptyset) \leq C_1 \exp(-C_2 n). \quad (21)$$

Indeed as in (3), given $H_n^a(\mathbf{u},\mathbf{v})$ and $H_n^b(\mathbf{u},\mathbf{v})$ property (c) of history regions allows us to define for every $n \geq 0$,

$$J_n^a(\mathbf{u}) := \inf\{m : U_{g_n^a(\mathbf{u})_m}^a < p\} \text{ and } J_n^b(\mathbf{v}) := \inf\{m : U_{g_n^b(\mathbf{v})_m}^b < p\}$$

as independent geometric random variables, each with success probability p. Let $\{G_n : n \geq 0\}$ be i.i.d. family of geometric random variables each with success probability p and, also independent of $\{U_{\mathbf{x}}^a : \mathbf{x} \in \mathbb{Z}^4\}$ and $\{U_{\mathbf{x}}^b : \mathbf{x} \in \mathbb{Z}^4\}$. Construct a sequence of random variables $J_0^{ind}, J_1^{ind}, J_2^{ind}, \ldots$, where for $n \geq 0$

$$J_n^{ind} = \begin{cases} \max\{J_n^a(\mathbf{u}), J_n^b(\mathbf{v})\} \text{ if } g_n^a(\mathbf{u})(4) = g_n^b(\mathbf{v})(4) \\ \max\{J_n^a(\mathbf{u}), G_n\} \text{ if } g_n^a(\mathbf{u})(4) < g_n^b(\mathbf{v})(4) \\ \max\{J_n^b(\mathbf{v}), G_n\} \text{ if } g_n^a(\mathbf{u})(4) > g_n^b(\mathbf{v})(4). \end{cases}$$

As in (4) we have for all m

$$\mathbb{P}(J_{n+1}^{ind} \leq m | g_n(\mathbf{u}), g_n(\mathbf{v}), H_n^a(\mathbf{u},\mathbf{v}), H_n^b(\mathbf{u},\mathbf{v})) = \mathbb{P}(\max\{G_1, G_2\} \leq m). \quad (22)$$

Now consider the Markov process $\{M_n^{ind}(\mathbf{u},\mathbf{v}), g_n^a(\mathbf{u}), H_n^a(\mathbf{u},\mathbf{v}), g_n^b(\mathbf{v}), H_n^b(\mathbf{u},\mathbf{v}) : n \geq 0\}$ where $M_n^{ind}(\mathbf{u},\mathbf{v})$ is defined in a similar fashion as $M_n(\mathbf{u},\mathbf{v})$ using J_n^{ind} instead of J_n in (5). Defining $\tau(M^{ind}(\mathbf{u},\mathbf{v})) := \inf\{n : M_n^{ind}(\mathbf{u},\mathbf{v}) = 0\}$, the coupling as in the first part of the proof of Proposition 2 gives $\tau^{ind}(\mathbf{u},\mathbf{v}) \leq \tau(M^{ind}(\mathbf{u},\mathbf{v}))$. Because of (22) we have

$$\tau(M^{ind}(\mathbf{u},\mathbf{v})) \overset{d}{=} \tau(M(\mathbf{u},\mathbf{v})) \tag{23}$$

and hence (21) is true.

Let

$$\begin{aligned} D_j^a &:= \{(\overline{g_n^a(\mathbf{u})} - \overline{g_{n-1}^a(\mathbf{u})}) : \tau_{j-1}^{ind}(\mathbf{u},\mathbf{v}) < n \leq \tau_j^{ind}(\mathbf{u},\mathbf{v})\} \\ D_j^b &:= \{(\overline{g_n^b(\mathbf{v})} - \overline{g_{n-1}^b(\mathbf{v})}) : \tau_{j-1}^{ind}(\mathbf{u},\mathbf{v}) < n \leq \tau_j^{ind}(\mathbf{u},\mathbf{v})\}; \end{aligned}$$

$\{(D_j^a, D_j^b) : j \geq 1\}$ is a collection of i.i.d. finite blocks of random variables with each of the blocks D_j^a and D_j^b being identically distributed for each $j \geq 1$. Thus, for $j \geq 0$

$$Z_j^{ind}(\mathbf{u},\mathbf{v}) := \overline{g^b_{\tau_j^{ind}(\mathbf{u},\mathbf{v})}(\mathbf{v})} - \overline{g^a_{\tau_j^{ind}(\mathbf{u},\mathbf{v})}(\mathbf{u})}$$

is a random walk on $\mathbb{Z}^3$ with zero mean and finite variance covariance matrix.

Now we construct the coupled process c. First we need some notation. Given a vertex $\mathbf{u} \in \mathbb{Z}^d$ let

$$\Delta_m(\mathbf{u}) := \begin{cases} \{\mathbf{w} \in \mathbb{Z}^d : \mathbf{u}(d) < \mathbf{w}(d) \leq m \text{ and } ||\mathbf{w} - \mathbf{u}||_1 \leq m - \mathbf{u}(d)\} & \text{if } m > \mathbf{u}(d) \\ \emptyset & \text{otherwise.} \end{cases}$$

Let $g_0^c(\mathbf{u}) := \mathbf{u}$, $g_0^c(\mathbf{v}) := \mathbf{v}$ and $H_0^c(\mathbf{u},\mathbf{v}) := \emptyset$.

Since the construction of the process is slightly different from that in Section 2, we give the details here. The reader should note that this construction yields two paths from $\mathbf{u}$ and $\mathbf{v}$ which are probabilistically equivalent to two paths from $\mathbf{u}$ and $\mathbf{v}$ in our original process. Define

$$\begin{aligned} & g_1^c(\mathbf{u}) := h^a(\mathbf{u}),\ g_1^c(\mathbf{v}) := \mathbf{v}, \\ & H_1^{c(a)}(\mathbf{u},\mathbf{v}) := \{(\mathbf{w}, U_{\mathbf{w}}^a) : \mathbf{w}(4) > \mathbf{v}(4),\ ||\mathbf{u} - \mathbf{w}||_1 \leq ||\mathbf{u} - g_1^c(\mathbf{u})||_1\}, \\ & H_1^{c(b)}(\mathbf{u},\mathbf{v}) := \emptyset, \\ & H_1^c(\mathbf{u},\mathbf{v}) := H_1^{c(a)}(\mathbf{u},\mathbf{v}) \cup H_1^{c(b)}(\mathbf{u},\mathbf{v}). \end{aligned}$$

Having obtained $g_k^c(\mathbf{u}), g_k^c(\mathbf{v})$ and the history sets $H_k^{c(a)}(\mathbf{u},\mathbf{v}), H_k^{c(b)}(\mathbf{u},\mathbf{v}), H_k^c(\mathbf{u},\mathbf{v})$ for all $0 \leq k \leq n$, we define recursively for the three cases


$H_1^{c(a)}(\mathbf{u},\mathbf{v})$
$g_1^c(\mathbf{u})$
$\mathbf{u}$
$\mathbf{v} = g_1^c(\mathbf{v})$
$H_4^{c(a)}(\mathbf{u},\mathbf{v})$
$g_4^c(\mathbf{u})$
$H_4^{c(b)}(\mathbf{u},\mathbf{v})$
$g_4^c(\mathbf{v})$
$\mathbf{u}$
$\mathbf{v}$
$H_2^{c(a)}(\mathbf{u},\mathbf{v})$
$g_2^c(\mathbf{u})$
$H_2^{c(b)}(\mathbf{u},\mathbf{v})$
$\mathbf{u}$
$g_2^c(\mathbf{v}) = \mathbf{v}$
$H_3^{c(a)}(\mathbf{u},\mathbf{v})$
$g_3^c(\mathbf{u})$
$H_3^{c(b)}(\mathbf{u},\mathbf{v})$
$\mathbf{u}$
$g_3^c(\mathbf{v}) = \mathbf{v}$

Figure 9: Construction of the coupled c process from $\mathbf{u}$ and $\mathbf{v}$. The darker shaded regions represent $H_n^{c(a)}(\mathbf{u},\mathbf{v})$ and the lighter shaded regions represent $H_n^{c(b)}(\mathbf{u},\mathbf{v})$.

(i) If $g_n^c(\mathbf{u})(4) = g_n^c(\mathbf{v})(4)$

(ii) If $g_n^c(\mathbf{u})(4) > g_n^c(\mathbf{v})(4)$, $g_{n-1}^c(\mathbf{v}) = g_n^c(\mathbf{v})$ and $g_{n-1}^c(\mathbf{u}) = g_n^c(\mathbf{u})$

(iii) If $g_n^c(\mathbf{u})(4) > g_n^c(\mathbf{v})(4)$ and $g_{n-1}^c(\mathbf{v}) \neq g_n^c(\mathbf{v})$ or $g_{n-1}^c(\mathbf{u}) \neq g_n^c(\mathbf{u})$.

In cases (i) and (ii), we set

$$\begin{aligned}
&g_{n+1}^c(\mathbf{u}) := h^a(g_n(\mathbf{u})), \quad g_{n+1}^c(\mathbf{v}) := g_n^c(\mathbf{v}),\\
&H_{n+1}^{c(a)}(\mathbf{u},\mathbf{v}) := H_n^{c(a)}(\mathbf{u},\mathbf{v}) \cup \{(\mathbf{w}, U_{\mathbf{w}}^a) : \mathbf{w}(4) > \max\{g_n^c(\mathbf{v})(4), g_n^c(\mathbf{u})(4)\} \text{ and}\\
&\qquad\qquad ||g_n^c(\mathbf{u}) - \mathbf{w}||_1 \leq ||g_n^c(\mathbf{u}) - g_{n+1}^c(\mathbf{u})||_1\},\\
&H_{n+1}^{c(b)}(\mathbf{u},\mathbf{v}) := H_n^{c(b)}(\mathbf{u},\mathbf{v}),\\
&H_{n+1}^{c}(\mathbf{u},\mathbf{v}) := H_{n+1}^{c(a)}(\mathbf{u},\mathbf{v}) \cup H_{n+1}^{c(b)}(\mathbf{u},\mathbf{v}).
\end{aligned}$$

For case (iii), consider

$$\begin{aligned}
\Delta_m^a(g_n^c(\mathbf{v})) &:= \Delta_m(g_n^c(\mathbf{v})) \cap \{\mathbf{w} : (\mathbf{w}, U_{\mathbf{w}}^a) \in H_n^{c(a)}(\mathbf{u},\mathbf{v})\}\\
\Delta_m^{b\backslash a}(g_n^c(\mathbf{v})) &:= \Delta_m(g_n^c(\mathbf{v})) \setminus \Delta_m^a(g_n^c(\mathbf{v}))\\
\Lambda_m^c(g_n^c(\mathbf{v})) &:= \{(\mathbf{w}, U_{\mathbf{w}}^a) : \mathbf{w} \in \Delta_m^a(g_n^c(\mathbf{v}))\} \cup \{(\mathbf{w}, U_{\mathbf{w}}^b) : \mathbf{w} \in \Delta_m^{b\backslash a}(g_n^c(\mathbf{v}))\}.
\end{aligned}$$

Define

$$m_{n+1} := \inf\{m : \text{ there exists } (\mathbf{w}, x) \in \Lambda_m^c(g_n^c(\mathbf{v})) \text{ with } x \leq p\}$$

$$\mathbf{w}_{n+1} := \begin{cases} \mathbf{z} & \text{if } m_{n+1} \leq g_n^c(\mathbf{u})(4) \\ \mathbf{0} & \text{otherwise,} \end{cases}$$

where $(\mathbf{z}, x) \in \Lambda^c_{m_{n+1}}(g_n^c(\mathbf{v}))$ and $x \leq y$ for all $(\mathbf{w}, y) \in \Lambda^c_{m_{n+1}}(g_n^c(\mathbf{v}))$.

For case (iii) we need to consider two situations.

If $m_{n+1} \leq g_n^c(\mathbf{u})(4)$

$$\begin{aligned} g^c_{n+1}(\mathbf{v}) &:= \mathbf{w}_{n+1} \\ g^c_{n+1}(\mathbf{u}) &:= g^c_n(\mathbf{u}) \\ H^{c(b)}_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(b)}_n(\mathbf{u},\mathbf{v}) \cup \{(\mathbf{w}, U^b_{\mathbf{w}}) : \mathbf{w} \in \Delta^{b\setminus a}_{m_{n+1}}(g^c_n(\mathbf{v}))\} \\ &\quad \cap \{(\mathbf{w}, x) : 0 \leq x \leq 1,\ \mathbf{w}(4) > g^c_{n+1}(\mathbf{v})(4)\} \\ H^{c(a)}_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(a)}_n(\mathbf{u},\mathbf{v}) \cap \{(\mathbf{w}, x) : 0 \leq x \leq 1,\ \mathbf{w}(4) > g^c_{n+1}(\mathbf{v})(4)\} \\ H^c_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(a)}_{n+1}(\mathbf{u},\mathbf{v}) \cup H^{c(b)}_{n+1}(\mathbf{u},\mathbf{v}) \end{aligned}$$

If $m_{n+1} > g_n^c(\mathbf{u})(4)$

$$\begin{aligned} g^c_{n+1}(\mathbf{v}) &:= g^c_n(\mathbf{v}) \\ g^c_{n+1}(\mathbf{u}) &:= g^c_n(\mathbf{u}) \\ H^{c(b)}_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(b)}_n(\mathbf{u},\mathbf{v}) \cup \{(\mathbf{w}, U^b_{\mathbf{w}}) : \mathbf{w} \in \Delta^{b\setminus a}_{g^c_n(\mathbf{u})(4)}(g^c_n(\mathbf{v}))\} \\ H^{c(a)}_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(a)}_n(\mathbf{u},\mathbf{v}) \\ H^c_{n+1}(\mathbf{u},\mathbf{v}) &:= H^{c(a)}_{n+1}(\mathbf{u},\mathbf{v}) \cup H^{c(b)}_{n+1}(\mathbf{u},\mathbf{v}) \end{aligned}$$

Set $\tau^c_0(\mathbf{u},\mathbf{v}) := 0$, and for $j \geq 1$, let

$$\tau^c_j(\mathbf{u},\mathbf{v}) := \inf\{n > \tau^c_{j-1}(\mathbf{u},\mathbf{v}) : H^c_n(\mathbf{u},\mathbf{v}) = \emptyset\},$$

denote the number of steps required for j th regeneration of the c process.

Define a $\mathbb{Z}^3$ valued process as,

$$Z^c_j(\mathbf{u},\mathbf{v}) := \bar{g}^c_{\tau^c_j(\mathbf{u},\mathbf{v})}(\mathbf{v}) - \bar{g}^c_{\tau^c_j(\mathbf{u},\mathbf{v})}(\mathbf{u}),$$

which is identically distributed with the process $\{Z_j(\mathbf{u},\mathbf{v}); j \geq 0\}$ defined in (16).

Consider the event that till the regeneration step $\tau^c_1(\mathbf{u},\mathbf{v})$, the set of vertices explored to obtain the path $\{g^c_n(\mathbf{v}) : 0 \leq n \leq \tau^c_1(\mathbf{u},\mathbf{v})\}$ is disjoint from the set of vertices explored to obtain the path $\{g^c_n(\mathbf{u}) : 0 \leq n \leq \tau^c_1(\mathbf{u},\mathbf{v})\}$. On this event $g^c_{\tau^c_1(\mathbf{u},\mathbf{v})}(\mathbf{v})$ depends only on the $\{U^b_{\mathbf{x}}\}$ values in the explored region and $g^b_{\tau^{ab}_1(\mathbf{u},\mathbf{v})}(\mathbf{v})$ and $g^a_{\tau^{ab}_1(\mathbf{u},\mathbf{v})}(\mathbf{u})$ are the same as $g^c_{\tau^c_1(\mathbf{u},\mathbf{v})}(\mathbf{v})$ and $g^c_{\tau^c_1(\mathbf{u},\mathbf{v})}(\mathbf{u})$ respectively. Fix $j \geq 1$, and let

$$W^{ab}_j(\mathbf{u},\mathbf{v}) := \sum_{i=\tau^{ab}_{j-1}(\mathbf{u},\mathbf{v})+1}^{\tau^{ab}_j(\mathbf{u},\mathbf{v})} \{||g^a_i(\mathbf{u}) - g^a_{i-1}(\mathbf{u})||_1 + ||g^b_i(\mathbf{v}) - g^b_{i-1}(\mathbf{v})||_1\}.$$

From (23), as in (14) we have, for any $l > 0$,

$$\mathbb{P}(W_j^{ab}(\mathbf{u}, \mathbf{v}) > l) \leq \mathbb{P}(\Gamma \geq l) \leq D_1 \exp -(D_2 l),$$

for positive constants D_1 and D_2; and hence

$$\mathbb{P}(Z_1^{ab}(\mathbf{u}, \mathbf{v}) = Z_1^{c}(\mathbf{u}, \mathbf{v})) \geq 1 - D_1 \exp(-D_2 \frac{||v||_2}{4}). \tag{24}$$

Given $\epsilon > 0$ define the event

$$\begin{aligned} A_{n,\epsilon}(\mathbf{u}, \mathbf{v}) := \{ & g^c_{\tau_{n^4}(\mathbf{u},\mathbf{v})}(\mathbf{v}) \in g^c_{\tau_{n^4}(\mathbf{u},\mathbf{v})}(\mathbf{u}) + (\triangle_{n^{2(1+\epsilon)}} \setminus \triangle_{n^{2(1-\epsilon)}}), \\ & g^c_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{v}) \neq g^c_{\tau_j(\mathbf{u},\mathbf{v})}(\mathbf{u}) \text{ for all } j = 1, \ldots, n^4\}, \end{aligned}$$

where $\triangle_n := \{\mathbf{x} \in Z^3 : ||\mathbf{x}||_1 \leq n\}$.

Lemma 1 *For $0 < \epsilon < \frac{1}{3}$, there exist constants $C, \beta > 0$ and $n_0 \geq 1$ such that, for all $n \geq n_0$,*

$$\inf_{\mathbf{v} \in \mathbf{u} + (\triangle_{n^{(1+\epsilon)}} \setminus \triangle_{n^{(1-\epsilon)}})} \mathbb{P}(A_{n,\epsilon}(\mathbf{u}, \mathbf{v})) \geq 1 - Cn^{-\beta}$$

Lemma 1 and its proof are similar to that of Lemma 3.2 of [GRS04]. Here we define an independent version of the event $A_{n,\epsilon}(\mathbf{u}, \mathbf{v})$ as,

$$\begin{aligned} B_{n,\epsilon}(\mathbf{u}, \mathbf{v}) := \{ & g^b_{\tau^{ab}_{n^4}(\mathbf{u},\mathbf{v})}(\mathbf{v}) \in g^a_{\tau^{ab}_{n^4}(\mathbf{u},\mathbf{v})}(\mathbf{u}) + (\triangle_{n^{2(1+\epsilon)}} \setminus \triangle_{n^{2(1-\epsilon)}}), \\ & ||g^b_{\tau^{ab}_j(\mathbf{u},\mathbf{v})}(\mathbf{v}) - g^a_{\tau^{ab}_j(\mathbf{u},\mathbf{v})}(\mathbf{u})||_1 \geq K \log_n \text{ for all } j = 1, \ldots, n^4\}, \end{aligned}$$

Similar calculations as in the proof of Lemma 3.2 of [GRS04] give us,

$$\inf_{\mathbf{v} \in \mathbf{u} + (\triangle_{n^{(1+\epsilon)}} \setminus \triangle_{n^{(1-\epsilon)}})} 1 - \mathbb{P}(B_{n,\epsilon}(\mathbf{u}, \mathbf{v})) < C_3 n^{-\alpha}, \tag{25}$$

which is a version of equation (19) of Lemma 3.2 of [GRS04]. The only difference is that our random walk is not isotropic. This difference is unimportant, since it has all the necessary properties like symmetry and finite covariance matrix. We also have (24), which is version of equation (17) of Lemma 3.2 of [GRS04]. From (24) and from (25), similar calculations as [GRS04] give us Lemma 1. As mentioned in Section 3.3 in the proof of Theorem 2.1 of [GRS04], from Lemma 1 it follows that $\mathbb{P}(G$ is disconnected $) > 0$.

Now by the inherent ergodicity of the process, we have that

$$\mathbb{P}(G \text{ is disconnected }) = 1.$$

A similar argument along with the ergodicity of the random graph, may further be used to establish that for any $k \geq 1$

$$\mathbb{P}(G \text{ has atleast } k \text{ trees }) = 1,$$

and thus

$$\mathbb{P}(G \text{ has infinitely many trees }) = 1.$$

4 Brownian Web

In this section we prove Theorem 2. We begin by recalling from [FINR04] and [NRS05] the relevant details of the Brownian web. Let $\mathbb{R}^2_c$ denote the completion of the space time plane $\mathbb{R}^2$ with respect to the metric

$$\rho((x_1,t_1),(x_2,t_2)) = |\tanh(t_1) - \tanh(t_2)| \vee |\frac{\tanh(x_1)}{1+|t_1|} - \frac{\tanh(x_2)}{1+|t_2|}|.$$

Let Π be the space of all paths in $\mathbb{R}^2_c$ of the form $\{(\pi(t),t) : t \geq \sigma_\pi\}$ where $\sigma_\pi \in [-\infty,\infty]$ denotes the starting time for path π. The following metric, for $\pi_1, \pi_2 \in \Pi$

$$d(\pi_1,\pi_2) = |\tanh(\sigma_{\pi_1}) - \tanh(\sigma_{\pi_2})| \vee \sup_{t \geq \sigma_{\pi_1} \wedge \sigma_{\pi_2}} |\frac{\tanh(\pi_1(t \vee \sigma_{\pi_1}))}{1+|t_1|} - \frac{\tanh(\pi_2(t \vee \sigma_{\pi_2}))}{1+|t_2|}|$$

makes Π a complete, separable metric space. Convergence in this metric can be described as locally uniform convergence of paths as well as convergence of starting times. Let $\mathcal{H}$ be the space of compact subsets of (Π, d), equipped with the Hausdorff metric $d_{\mathcal{H}}$ given by,

$$d_{\mathcal{H}}(K_1,K_2) = \sup_{\pi_1 \in K_1} \inf_{\pi_2 \in K_2} d(\pi_1,\pi_2) \vee \sup_{\pi_2 \in K_2} \inf_{\pi_1 \in K_1} d(\pi_1,\pi_2).$$

The space $(\mathcal{H}, d_{\mathcal{H}})$ is a complete separable metric space. Let $B_{\mathcal{H}}$ be the Borel $\sigma-$algebra on the metric space $(\mathcal{H}, d_{\mathcal{H}})$. The Brownian web $\mathcal{W}$ is an $(\mathcal{H}, B_{\mathcal{H}})$ valued random variable.

We now recall from [FINR04, Theorem 2.1] the following characterization of the Brownian web $\mathcal{W}$.

Theorem 3 *There exists an $(\mathcal{H}, B_{\mathcal{H}})$ valued random variable $\mathcal{W}$, called the standard Brownian web, whose distribution is uniquely determined by the following properties:*

(i) For each deterministic $z \in \mathbb{R}^2$, almost surely there is a unique path $\pi_z \in \mathcal{W}(z)$.

(ii) For any finite deterministic set of points $z_1, \ldots, z_k \in \mathbb{R}^2$, the collection $(\pi_{z_1}, \ldots, \pi_{z_k})$ is distributed as coalescing Brownian motions.

(iii) For any deterministic countable dense subset $\mathcal{D} \subset \mathbb{R}^2$, almost surely, $\mathcal{W}$ is the closure of $\{\pi_z : z \in \mathcal{D}\}$ in (Π, d).

For a compact set of paths $K \in \mathcal{H}$ and for $t > 0$ and $t_0, a, b \in \mathbb{R}$ with $a < b$, let

$$\eta_K(t_0, t; a, b) := \#\{\pi(t_0 + t) : \pi \in K \text{ with } \sigma_\pi \le t_0 \text{ and } \pi(t_0) \in [a, b]\},$$

be the $\{0, 1, 2, \ldots, \infty\}$ valued random variable giving the number of distinct points on $\mathbb{R} \times \{t_0 + t\}$ intersected by some path in K which also intersects $[a, b] \times \{t_0\}$. Consider the dual family of counting random variables

$$\begin{aligned}\hat{\eta}_X(t_0, t; a, b) :=& \#\{x \in (a, b) : \text{ there exists a path in } X \text{ intersecting both} \\ & \mathbb{R} \times \{t_0\} \text{ and } (x, t_0 + t)\}\end{aligned}$$

We prove Theorem 2 by verifying the convergence criteria for the Brownian web for $(\mathcal{H}, B_{\mathcal{H}})$ valued random variables with non-crossing paths, proposed by [FINR04] and [NRS05], as the following:

Theorem 4 *Let $(\chi_n)_{n\in\mathbb{N}}$ be a sequence of $(\mathcal{H}, B_{\mathcal{H}})$ valued random variables with non-crossing paths. If the following conditions are satisfied then χ_n converges in distribution to the standard Brownian web $\mathcal{W}$.*

- (I_1) *Let $\mathcal{D}$ be a countable deterministic dense subset of $\mathbb{R}^2$. There exists $\theta_n^{\mathbf{x}} \in \chi_n$ for $\mathbf{x} \in \mathcal{D}$ such that for any finite deterministic collection $\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_m \in \mathcal{D}$, $({\theta_n}^{\mathbf{x}_1}, \ldots, {\theta_n}^{\mathbf{x}_m})$ converges in distribution as $n \to \infty$ to coalescing Brownian motions starting from $(\mathbf{x}_1, \mathbf{x}_2, \ldots, \mathbf{x}_m)$.*
- (B_1) *For all $t > 0, \limsup_{n\to\infty} sup_{(a,t_0)\in\mathbb{R}^2} \mathbb{P}(\eta_{\chi_n}(t_0, t; a, a + \epsilon) \ge 2) \to 0$ as $\epsilon \to 0$*
- (E_1) *If χ is any subsequential limit of χ_n, then for all $t_0, \epsilon, a, b \in \mathbb{R}$ with $\epsilon > 0$ and $a < b$, $\mathbb{E}[\hat{\eta}_\chi(t_0, t; a, b)] \le \mathbb{E}[\hat{\eta}_W(t_0, t; a, b)] = \frac{b-a}{\sqrt{t\pi}}$.*

4.1 Proof of condition (I_1)

For $\chi_n(\gamma, \sigma)$ as in (2) to show (I_1) we need to control the tail of the distribution of the coalescing time of two paths starting at the same time at unit distance apart.

Lemma 2 *For $(0, 0), (1, 0) \in V$, let $\nu = \inf\{j : g_{\tau_j((0,0),(1,0))}(0, 0) = g_{\tau_j((0,0),(1,0))}(1, 0)\}$. Then there exists a constant $C > 0$ such that we have*

$$\mathbb{P}(T_\nu((0, 0), (1, 0)) > t) \le \frac{C}{\sqrt{t}}.$$

Proof: Consider the nonnegative martingale, $\{Z_j((0, 0), (1, 0)), \mathcal{F}_{T_j((0,0),(1,0))}\}$, and we proceed in the same way as in Theorem 3 in [CFD09]. For $m \ge 1$, we want to find an upperbound for $\mathbb{P}(Z_{j+1}((0, 0), (1, 0)) = m|Z_j((0, 0), (1, 0)) = m)$.



Figure 10:

Consider the situation as in the Figure 10 where the vertices $(0,0)$, $(0,2)$, $(m,0)$, $(m+1,1)$, $(m+1,2)$ are open and all other nearby vertices closed, so as to give paths as shown in the figure.

We thus clearly have

$$\begin{aligned}&\mathbb{P}(Z_{j+1}((0,0),(1,0)) = m | Z_j((0,0),(1,0)) = m)\\ \leq\ & 1 - \mathbb{P}(Z_1((0,0),(m,0)) = m+1 | Z_0((0,0),(m,0)) = m)\\ \leq\ & 1-(1-p)^6p^3.\end{aligned}$$

This upper bound is independent of m and for $0 < p < 1$, it is strictly smaller than 1. Now the same proof as in Lemma 1 of [CFD09], gives us a bound on the number of regeneration steps before coalescing,

$$\mathbb{P}(\nu > t) \leq \frac{c_1}{\sqrt{t}} \text{ for some } c_1 > 0.$$

Since the time elapsed between any two consecutive regeneration steps is stochastically dominated by Γ, the rest of the proof follows easily. ☐

Proposition 4 *Let* $(\mathbf{x}_1, \dots, \mathbf{x}_m) \in \mathbb{R}^2$ *be such that* $\mathbf{x}_1(2) \leq \mathbf{x}_2(2) \leq \cdots \leq \mathbf{x}_m(2)$ *with* $\mathbf{x}_{i-1}(1) < \mathbf{x}_i(1)$ *whenever* $\mathbf{x}_{i-1}(2) = \mathbf{x}_i(2)$*. For* $\sigma, \gamma > 0$ *let*

$$Z_n^{(i)}(\gamma, \sigma) = \{Z_n^{(i)}(t) := \frac{1}{n\sigma}\pi^{([\mathbf{x}_i(1)n\sigma],[\mathbf{x}_i(2)n^2\gamma])}([\gamma n^2 t]); t \geq \mathbf{x}_i(2)\}.$$

Then there exist $\gamma := \gamma(p)$ *and* $\sigma := \sigma(p)$ *such that, as* $n \to \infty$*, we have*

$$\{Z_n^{(i)}(\gamma, \sigma), i = 1, 2, \dots, m\} \text{ converges weakly to } \{W^{(i)}, i = 1, 2, \dots, m\},$$

where $W^{(i)}$*'s are standard coalescing Brownian motions starting from* $\mathbf{x}_i$*'s.*

Proof: We prove this by induction. Start with $m = 1$ and without loss of generality assume that $\mathbf{x}_1$ is such that $\mathbf{x}_1(2) = 0$. Note that $\{g_{\tau_n(\mathbf{x}_1)}(\mathbf{x}_1)(1) : n \geq 1\}$ is a symmetric random walk with increments having moments of all order. Let $\tilde{\pi}^{\mathbf{x}_1}$ denote the path obtained through linear interpolation of the consecutive steps $\{g_{\tau_j(\mathbf{x}_1)}(\mathbf{x}_1), g_{\tau_{j+1}(\mathbf{x}_1)}(\mathbf{x}_1) : j \geq 0\}$. For $n \geq 1$ let

$$\tilde{Z}_n^{(1)}(\gamma, \sigma)(t) := \frac{1}{n\sigma}\tilde{\pi}^{\mathbf{x}_1^n}(\gamma n^2 t) \text{ for } t \geq 0$$

where $\mathbf{x}_1^n := ([n\mathbf{x}_1(1)\sigma], [n^2\mathbf{x}_1(2)\gamma])$. Now $\tilde{\pi}^{\mathbf{x}_1}$ can be regarded as a time change of the random walk $X(j) := \{g_{\tau_j(\mathbf{x}_1)}(\mathbf{x}_1)(1) : j \geq 0\}$ where the time change is given by $T(j) := T_j(\mathbf{x}_1)$, and $X(t)$ and $T(t)$ for non-integer t are defined by linearly interpolating between consecutive integer times. More precisely $\tilde{\pi}^{(\mathbf{x}_1)}(t) = X(T^{-1}(t))$. Hence we have

$$\tilde{Z}_n^{(1)}(\gamma, \sigma)(t) = n^{-1}\sigma^{-1}X(n^2\frac{1}{n^2}T^{-1}(n^2\gamma t)) \text{ for } t \geq 0$$

Note that $T^{-1}(n^2\gamma t)$ represents the number of renewals until the time $n^2\gamma t$. Hence by the renewal theorem $\frac{1}{n^2}T^{-1}(n^2\gamma t)$ converges in distribution to the linear function $g(t) := \frac{\gamma t}{\mathbb{E}(T_1(\mathbf{x}_1))}$, $t \geq 0$. Taking $\gamma_0 := \mathbb{E}(T_1(\mathbf{x}_1))$ as the expected regeneration time and σ_0 as the standard deviation of the random walk $X(n)$, from Donsker's invariance principle we have that $\tilde{Z}_n^{(1)}(\gamma_0, \sigma_0)$ converges weakly to the Brownian motion $W^{(1)}$ starting from $\mathbf{x}_1$.

Since,

$$\sup_{t \in [T_j(\mathbf{x}_1^n), T_{j+1}(\mathbf{x}_1^n))} |\pi^{\mathbf{x}_1^n}(t) - \tilde{\pi}^{\mathbf{x}_1^n}(t)| \leq W_j(\mathbf{x}_1^n)$$

where $W_j(\mathbf{x}_1^n)$ is as in (13), from (15) it follows that $Z_n^{(1)}(\gamma_0, \sigma_0)$ has the same weak limit as that of $\tilde{Z}_n^{(1)}(\gamma_0, \sigma_0)$. Hence the proposition holds for $m = 1$.

Now suppose $m \geq 2$ and consider first the case that $\mathbf{x}_1(2) = \mathbf{x}_2(2) = \cdots = \mathbf{x}_m(2) = 0$. We use ideas introduced in [CV11]. First with a little abuse of notation for $1 \leq i \leq m$ and for $n \geq 1$ let $\mathbf{x}_i^n := ([n\mathbf{x}_i(1)\sigma_0], [n^2\mathbf{x}_i(2)\gamma_0])$ and for ease in writing let $\pi_1, \dots, \pi_m$ denote the paths $\pi^{\mathbf{x}_1^n}, \pi^{\mathbf{x}_2^n}, \dots, \pi^{\mathbf{x}_m^n}$. The idea of the proof is to modify the paths $\pi_1, \dots, \pi_m$ in such a manner that the modified path π_m is independent of the remaining paths. Thus enabling us to conclude from the induction hypothesis that the first $m-1$ paths converge to coalescing Brownian motions, while the last being independent also coalesces with these coalescing Brownian motions. Our task would then be to show that this modification can be done with a probability which goes to 1 as $n \to \infty$.

Before we embark on the proof we need a notation. The exploration box for the j th regeneration of the path $\{h^n(\mathbf{u}), n \geq 0)\}$ is denoted by

$$D_j(\mathbf{u}) := \pi^{\mathbf{u}}(T_j(\mathbf{u})) + [-W_j(\mathbf{u}), W_j(\mathbf{u})] \times [T_j(\mathbf{u}), T_{j+1}(\mathbf{u})]. \tag{26}$$

We start with two independent collections, a green collection $\{U_{\mathbf{x}}^g : \mathbf{x} \in \mathbb{Z}^2\}$ and a red collection $\{U_{\mathbf{x}}^r : \mathbf{x} \in \mathbb{Z}^2\}$ of i.i.d. $U(0,1)$ random variables which are also independent of the original collection $\{U_{\mathbf{x}} : \mathbf{x} \in \mathbb{Z}^2\}$.

For $1 \leq i \leq m$ let

$$\sigma_i = \sigma_i^{(n)} := \inf\{j : W_j(\mathbf{x}_i^n) \geq n^\alpha\},$$

where $\alpha > 0$ is a suitable constant to be specified later, i.e. σ_i is the first regeneration step such that the width of the explored region to obtain the next regeneration is suitably large. (This is represented by the box for the path π_i in Figure 11.)



Figure 11: Construction of the original path π_i and the modified green path π_i^g.

For $1 \leq i \leq m-1$, let π_i^g denote the path obtained from the collection $\{U_{\mathbf{x}} : \mathbf{x}(2) \leq T_{\sigma_i}(\mathbf{x}_i^n)\} \cup \{U_{\mathbf{x}}^g : \mathbf{x}(2) > T_{\sigma_i}(\mathbf{x}_i^n)\}$. Thus π_i^g, the green path, coincides with the original path π_i until time $T_{\sigma_i}(\mathbf{x}_i^n)$, and thereafter it is independent of the original path.

Let $D_0^g(i), D_1^g(i), \ldots$ be the exploration boxes of widths $W_0^g(i), W_1^g(i), \ldots$ respectively, obtained from the regeneration steps $\tau_0^g(i), \tau_1^g(i), \ldots$ of the path π_i^g. For $t \in \mathbb{N}$, let $r_t := \max\{x : (x,t) \in \cup_{i=1}^{m-1} \cup_{j \geq 0} D_j^g(i)\}$ denote the rightmost boundary point at a height t of the exploration boxes of the first $m-1$ green paths and $l_t := \min\{x : (x,t) \in \cup_{j\geq 0} D_j(m)\}$ denote the leftmost boundary point at a height t of the exploration boxes $D_0(m), D_1(m), \ldots$ of the path π_m obtained from the original process. Let $t^* = \min\{t : |r_t - l_t| \leq 2n^\alpha\}$, and let

$$\mu_m = \mu_m^n := \inf\{j : \text{ there exists } (l_{t^*}, t^*) \in D_j(m)\},$$

i.e., $D_{\mu_m}(m)$ is the first exploration box of the path π_m which comes closer than a distance $2n^\alpha$ from some exploration box of the first $m-1$ green paths.

Let π_m^r denote the red path obtained from the collection $\{U_{\mathbf{x}} : \mathbf{x}(2) \leq \min\{T_{\mu_m}(\mathbf{x}_m^n), T_{\sigma_m}(\mathbf{x}_m^n)\} \cup \{U_{\mathbf{x}}^r : \mathbf{x}(2) > \min\{T_{\mu_m}(\mathbf{x}_m^n), T_{\sigma_m}(\mathbf{x}_m^n)\}\}$

and if π_m^r intersects any of the $m-1$ paths $\pi_1^g, \dots, \pi_{m-1}^g$ at a lattice point then from that point onwards it coalesces with the intersected green path. Thus π_m^r, the red path, coincides with the original path π_m until time $\min\{T_{\mu_m}(\mathbf{x}_m^n), T_{\sigma_m}(\mathbf{x}_m^n)\}$, and thereafter it is independent of the original path as well as the first $m-1$ green paths till coalescence.

Now let

$$B_{n,s} := \{T_{\sigma_i}(\mathbf{x}_i^n) > sn^2 : \text{ for all } 1 \le i \le m\},$$

represent the event that until time sn^2 each of the the green paths coincide with its corresponding original path and that no exploration box of the path π_m has width larger than n^α. Clearly, from

$$\begin{aligned}\mathbb{P}(B_{n,s}^c) &\le m\mathbb{P}(\cup_{j=1}^{sn^2}(W_j(1) \ge n^\alpha)) \\ &\le msn^2\mathbb{P}(W_1(1) \ge n^\alpha) \\ &\le msn^2\mathbb{P}(\Gamma \ge n^\alpha).\end{aligned}$$

and hence from (11) it follows that for any $\alpha > 0$ we have $\mathbb{P}(B_{n,s}^c) \to 0$ as $n \to \infty$.

Let

$$\begin{aligned}&T^o = T^{o,n} := \inf\{t : \pi_i(t) = \pi_m(t) \text{ for some } 1 \le i \le m-1\} \text{ and} \\ &T^r = T^{r,n} := \inf\{t : \pi_i(t) = \pi_m^r(t) \text{ for some } 1 \le i \le m-1\}.\end{aligned}$$

Taking $A_{n,s} := \{T_{\mu_m}(\mathbf{x}_m^n) \ge sn^2\}$, as in the proof of Proposition 3.7 of [CV11], it suffices to show that, as $n \to \infty$,

$$\mathbb{P}((T^o > T_{\mu_m}(\mathbf{x}_m^n) + n^{\frac{7}{4}}) \cap B_{n,s} \cap A_{n,s}^c) \to 0 \text{ and} \tag{27}$$

$$\mathbb{P}((T^r > T_{\mu_m}(\mathbf{x}_m^n) + n^{\frac{7}{4}}) \cap B_{n,s} \cap A_{n,s}^c) \to 0. \tag{28}$$

On the event $B_{n,s} \cap A_{n,s}^c$, the exploration box $D_{\mu_m}(m)$ lies at a distance smaller that $2n^\alpha$ from the exploration box $D_j(i)$ for some $1 \le i \le m-1$ and some $j \ge 0$. Let $t_0 := \max\{t : (x,t) \in D_j(i) \cup D_{\mu_m}(m)\}$ and start with two vertices $\mathbf{x}_r$ and $\mathbf{x}_l$ on the line $Y = t_0$ where

$$\mathbf{x}_r = \mathbf{x}_r^n := (\pi_m(T_{\mu_m+1}(\mathbf{x}_m^n)), t_0) + (2n^\gamma, 0) \text{ and } \mathbf{x}_l = \mathbf{x}_l^n := (\pi_i(T_{j+1}(\mathbf{x}_i^n)), t_0) - (2n^\gamma, 0).$$

Consider two paths π_l and π_r starting from $\mathbf{x}_l$ and $\mathbf{x}_r$ respectively using the original collection of uniform random variables $\{U_\mathbf{x} : \mathbf{x} \in \mathbb{Z}^2\}$. Let $T^{l,r} := \inf\{t : t \ge t_0, \pi_l(t) = \pi_r(t)\}$ denote the coalescing time for π_l and π_r. Note that for $T^{l,r}$ we have nice tail estimates from Lemma 2. On the event

$$F := \{\max\{W_j(i), W_{j+1}(i), W_{\mu_m}(m), W_{\mu_m+1}(m)\} \le n^\alpha\},$$



Figure 12: On the event $B_{n,s} \cap (A_{n,s})^c \cap F$, for $t \geq T_{\mu_m}(\mathbf{x}^n_m)$ and for some $1 \leq i \leq m-1$ both the original paths $\pi(i)$ and $\pi(m)$ get sandwiched between π_l and π_r.

both $\pi_i(t_0)$ and $\pi_m(t_0)$ lie between $\mathbf{x}_l$ and $\mathbf{x}_r$; and neither $\mathbf{x}_l$ nor $\mathbf{x}_r$ lie in the exploration boxes of π_i and π_m. Hence, on F we have $T^{0,n} \leq T^{l,r}$ (See Figure 12).

For $\alpha < \gamma < \frac{7}{8}$, $p \in (0,1)$ and from (15) we have,

$$\begin{aligned}
&\mathbb{P}(\{T^o > T_{\mu_m}(\mathbf{x}^n_m) + n^{\frac{7}{4}}\} \cap B_{n,s} \cap (A_{n,s})^c) \\
&\leq \mathbb{P}(\{T^o > T_{\mu_m}(\mathbf{x}^n_m) + n^{\frac{7}{4}}\} \cap B_{n,s} \cap F \cap (A_{n,s})^c) + \mathbb{P}(F^c) \\
&\leq \mathbb{P}(\{T^{l,r} > T_{\mu_m}(\mathbf{x}^n_m) + n^{\frac{7}{4}}\} \cap B_{n,s} \cap F \cap (A_{n,s})^c) + \mathbb{P}(F^c) \\
&\leq \frac{c_1 n^{\gamma}}{\sqrt{n^{\frac{7}{4}} - 2n^{\alpha}}} + \mathbb{P}(F^c) \\
&\to 0 \text{ as } n \to \infty.
\end{aligned}$$

This completes the proof for (27).

Let $T^{ind}_j(\mathbf{x}^n_m)$ denote the time taken for j th regeneration of the two independent processes $\{\pi_m(t) : t \geq T_{\mu_m}(\mathbf{x}^n_m)\}$ and $\{\pi^r_m(t) : t \geq T_{\mu_m}(\mathbf{x}^n_m)\}$ starting from $g_{\tau_{\mu_m}(m)}(\mathbf{x}^n_m)$. From our earlier discussion it follows that $\{\pi_m(T^{ind}_j(\mathbf{x}^n_m)) : j \geq 0\}$ and $\{\pi^r_m(T^{ind}_j(\mathbf{x}^n_m)) : j \geq 0\}$ are two independent recurrent random walks. Hence the coalescing time for the modified path π^r_m and the original path π_m is lesser than the time taken by the random walk $\{\pi^r_m(T^{ind}_j(\mathbf{x}^n_m)) - \pi_m(T^{ind}_j(\mathbf{x}^n_m)) : j \geq 0\}$ starting from 0 to hit 0. Since the above mentioned random walk has finite second order moment, by Proposition P3 of Spitzer [(2001),page 381], the proof for (28) follows. The remaining part of the proof for Proposition 4 is similar as in the Proposition 3.7 in [CV11].

We now consider the general case and proceed by induction on $m \geq 1$. We use ideas introduced in the proof of Part III in Theorem 4 of [CFD09]. For $m = 1$ there is nothing to prove. Suppose that for some $m \geq 2$ the result holds for $m - 1$. Let i_0 be the maximum of i's such that $\mathbf{x}_i(2) > \mathbf{x}_{i-1}(2)$. By the induction hypothesis

$$(\{Z_n^{(i)}(\gamma_0, \sigma_0)(t) : \mathbf{x}_i(2) \leq t \leq \mathbf{x}_{i_0}(2)\}, 1 \leq i < i_0) \text{ converges weakly to}$$

$$(\{W^{(i)}(t) : \mathbf{x}_i(2) \leq t \leq \mathbf{x}_{i_0}(2)\}, 1 \leq i < i_0),$$

and by Skorohod's representation theorem we can suppose that the convergence is almost sure. We first introduce some notations. For an $(\mathcal{H}, \mathcal{B}_{\mathcal{H}})$ valued random variable X, define X^{s^-} to be the subset of paths in X which start before or at time s and for $s \leq t$, define $X_T^{s^-,t}$ to be the set of paths in X^{s^-} truncated before time t. When $s = t$, we denote $X_T^{s^-,t}$ simply by X_T^s. Also let $X(t) \subset \mathbb{R}$ denote the set of values at time t of all paths in X. With a little abuse of notation $X(t)$ is also used to denote the set of points $(X(t), t) \subset \mathbb{R}^2$. For $1 \leq i < i_0$ let

$$\eta_i = \eta_i^n := \min\{j : T_j(\mathbf{x}_i^n) \geq \mathbf{x}_{i_0}^n(2)\} \text{ and } T_{max} = T_{max}^n := \max\{T_{\eta_i^n}(\mathbf{x}_i^n) : 1 \leq i < i_0\}.$$

Consider the random point sets

$$\begin{aligned} A_n &:= \{(\pi_i(T_{\eta_i}(\mathbf{x}_i^n)), T_{max}) : 1 \leq i < i_0\} \cup \{(\mathbf{x}_{i_0}^n(1), T_{max}), \ldots, (\mathbf{x}_m^n(1), T_{max})\}, \\ A &:= \{W^i : 1 \leq i < i_0\}(\mathbf{x}_{i_0}(2)) \cup \{\mathbf{x}_{i_0}, \mathbf{x}_{i_0+1}, \ldots, \mathbf{x}_m\}. \end{aligned}$$

All the points in $A_n \subset \mathbb{Z}^2$ are on the line $Y = T_{max}$ and we do not have any information about the uniform random variables $\{U_{\mathbf{x}} : \mathbf{x}(2) > T_{max}\}$. (See Figure 13.)

It is easy to see that under the diffusive scaling A_n converges to A almost surely as $n \to \infty$. Let χ^F denote a $(\mathcal{H}, \mathcal{B}_{\mathcal{H}})$ random variable consisting of paths starting from a random subset of $\mathbb{Z}^2$ distributed as F and let $\chi_n^F(\gamma_0, \sigma_0) := \{(\frac{\mathbf{x}(1)}{n\sigma_0}, \frac{\mathbf{x}(2)}{n^2\gamma_0}) : \mathbf{x} \in \chi^F\}$. By Lemma 6.5 of [NRS05] and by part 1 of Proposition 4, we have,

$$\chi_n^{A_n}(\gamma_0, \sigma_0) \text{ converges weakly to } W^A,$$

where W^A denote the process of coalescing Brownian motions starting from a random point set distributed as A. Let

$$\bar{\pi}_n(\gamma_0, \sigma_0) := \{(\frac{\mathbf{x}(1)}{n\sigma_0}, \frac{\mathbf{x}(2)}{n^2\gamma_0}) : \mathbf{x} \in \pi_i \text{ for some } 1 \leq i \leq m\}$$

denote a $(\mathcal{H}, B_{\mathcal{H}})$ valued random variable which consists of diffusively scaled original paths $\pi_1, \ldots, \pi_m$. We need to show that $\chi_n^{A_n}(\gamma_0, \sigma_0)$ and $\bar{\pi}_n(\gamma_0, \sigma_0)^{\mathbf{x}_{i_0}(2)_T}$



Figure 13: The construction of the original paths π_i for $1 \leq i \leq m$ and the set of paths in χ^{A_n} for $m = 2$. The two shaded vertices form the set A_n. The coalescing time between the paths in χ^{A_n} and π_i for $1 \leq i \leq m$ converges to 0 under diffusive scaling.

have the same weak limit. Fix a uniformly continuous bounded function $H : D([0,\infty])^m \to \mathbb{R}$. It suffices to show that

$$\lim_{n\to\infty} \mathbb{E}|[H(\chi_n^{A_n}(\gamma_0,\sigma_0)) - H(\bar{\pi}_n(\gamma_0,\sigma_0)_T^{\mathbf{x}_{i_0}(2)})]| = 0.$$

Fix $0 < \gamma_1 < 2$. Consider the event E_n where,

$$E_n := \{\chi_n^{A_n}(\gamma_0,\sigma_0)(t) = \bar{\pi}_n(\gamma_0,\sigma_0)_T^{\mathbf{x}_{i_0}(2)}(t) : t \geq \mathbf{x}_{i_0}(2) + n^{\gamma_1 - 2}\}.$$

We have

$$\begin{aligned}&\mathbb{E}|[H(\chi_n^{A_n}) - H(\bar{\pi}_n(\gamma_0,\sigma_0)_T^{\mathbf{x}_{i_0}(2)})]| \\ &\leq \mathbb{E}|[H(\chi_n^{A_n}) - H(\bar{\pi}_n(\gamma_0,\sigma_0)_T^{\mathbf{x}_{i_0}(2)})]1_{E_n}| + ||H||_\infty \mathbb{P}(E_n^c).\end{aligned}$$

By uniform continuity of H the first expectation converges to zero as $n \to \infty$. Using a similar sandwiching technique we can show that the paths in $\chi_n^{A_n}(\gamma_0,\sigma_0)$ coalesce with the paths in $\bar{\pi}_n(\gamma_0,\sigma_0)_T^{\mathbf{x}_{i_0}(2)}$ fast and infact $\mathbb{P}(E_n^c) \to 0$ as $n \to \infty$. This finally proves Proposition 4. ☐

In order to prove condition (E_1) we need the following lemma.

Lemma 3 *Suppose at time* 0 *paths* $\{g_j(z,0) : j \geq 0\}$ *start from* $(z,0)$ *for all* $z \in \mathbb{Z}$. *For* $t \geq 1$, *let* E_t *be the event that there exist* $j \in \mathbb{N}$ *and* $z \in \mathbb{Z}$ *such that* $g_j(z,0) = (0,t)$, *then*

$$\mathbb{P}(E_t) \leq \frac{C}{\sqrt{t}}$$

for some positive constant C *independent of everything else.*

Lemma 3 and its proof is similar to that of Lemma 2.7 in [NRS05].

Let Z_{t_0} be any subsequential limit of $\chi_n^{t_0^-}(\gamma_0, \sigma_0)$. Proceeding in the same way as in [NRS05] we first obtain that for any $\epsilon > 0$, $Z_{t_0}(t_0 + \epsilon)$ is locally finite almost surely. We have to show that $Z_{t_{0_T}}^{(t_0+\epsilon)}$, the set of paths in Z_{t_0} truncated before time $t_0 + \epsilon$, is distributed as $W^{Z_{t_0}(t_0+\epsilon)}$, i.e., coalescing Brownian motions starting from the random set $Z_{t_0}(t_0 + \epsilon) \subset \mathbb{R}^2$. It is enough to show that the finite dimensional distributions converge. Fix $-\infty < a < b < \infty$. Let

$$Z_{t_{0_T},(a,b)}^{(t_0+\epsilon)} := \{\pi : \pi \in Z_{t_{0_T}}^{(t_0+\epsilon)} \text{ and } \pi(t_0 + \epsilon) \in (a,b)\}$$

denote the set of paths in Z_{t_0} which pass through $(a,b) \times (t_0 + \epsilon)$ and let

$$Z_{t_0}^{\epsilon}(a,b) := Z_{t_{0_T},(a,b)}^{(t_0+\epsilon)}(t_0 + \epsilon) = \{\pi(t_0 + \epsilon) : \pi \in Z_{t_{0_T},(a,b)}^{(t_0+\epsilon)}\}.$$

It suffices to show that $Z_{t_{0_T},(a,b)}^{(t_0+\epsilon)}$ is distributed as $W^{Z_{t_0}^{\epsilon}(a,b)}$, i.e., coalescing Brownian motions starting from the random point set $Z_{t_0}^{\epsilon}(a,b) \subset \mathbb{R}^2$. As mentioned in [NRS05], the keypoint is to observe the paths as processes regenerating from a suitable (random) point set so that we can use Proposition 4 along with the Lemma 6.5 in [NRS05]. In case of our model consider the regeneration steps such that for the first time the vertical distance traversed in the rescaled lattice are greater than or equal to $t_0 + \epsilon$. Suppose $\bar{T}_{max}$ is the maximum vertical distance traversed. Consider the random point set consists of projection of regeneration steps on the line $Y = \bar{T}_{max}$. By what we have done earlier it follows that paths originating from this random point set and the original paths have the same weak limit $W^{Z_{t_0}^{\epsilon}(a,b)}$.
Acknowledgements: Kumarjit Saha is grateful to Indian Statistical Institute for a fellowship to pursue his Ph.D.

References

[A79] R. Arratia. Coalescing Brownian motions on the line. Ph.D. Thesis, University of Wisconsin, Madison, 1979.

[A81] R. Arratia. Coalescing Brownian motions and the voter model on $\mathbb{Z}$. Unpublished partial manuscript. Available from rarratia@math.usc.edu.

[BB07] F. Bacelli and C. Bordenave. The radial spanning tree of a Poisson point process. *Ann. Appl. Probab.* 17, 305–359, 2007.

[CFD09] C.F. Coletti, L.R.G. Fontes, and E.S. Dias. Scaling limit for a drainage network model. *J. Appl. Probab.* 46, 1184–1197, 2009.

[CV11] C.F. Coletti and G. Valle. Convergence to The Brownian Web for a generalization of the drainage network model. Preprint, arXiv:1109.3517v1, 2011.

[CT11] D. Coupier and V.C. Tran. The Directed Spanning Forest is almost surely a tree. Preprint, arXiv:1010.0773v2, 2011.

[FFW05] P.A. Ferrari, L.R.G. Fontes, and X.-Y. Wu. Two-dimensional Poisson Trees converge to the Brownian web. *Ann. Inst. H. Poincaré Probab. Statist.* 41, 851–858, 2005.

[FINR04] L.R.G. Fontes, M. Isopi, C.M. Newman, and K. Ravishankar. The Brownian web: characterization and convergence. *Ann. Probab.* 32(4), 2857–2883, 2004.

[FLT04] P.A. Ferrari, C. Landim, and H. Thorisson. Poisson trees, succession lines and coalescing random walks. *Ann. Inst. H. Poincaré Probab. Statist.* 40, 141–152, 2004.

[GRS04] S. Gangopadhyay, R. Roy, and A. Sarkar. Random oriented Trees: a Model of drainage networks. *Ann. Appl. Probab.* 14, 1242–1266, 2004.

[NRS05] C.M. Newman, K. Ravishankar, and R. Sun. Convergence of coalescing nonsimple random walks to the Brownian web. *Electron. J. Prob.* 10, 21–60, 2005.

[RR97] Rodriguez-Iturbe, I. and Rinaldo, A. (1997). Fractal river basins: chance and self-organization. Cambridge Univ. Press, New York.

[S67] Scheidegger, A. E. (1967). A stochastic model for drainage pattern into an intramontane trench. *Bull. Ass. Sci. Hydrol.* 12, 15-20.

[S01] Spitzer, F. (2001). Principles of Random Walk. Springer.

Rahul Roy, Kumarjit Saha and Anish Sarkar
Theoretical Statistics and Mathematics Unit
Indian Statistical Institute

7 S. J. S. Sansanwal Marg
New Delhi 110016, India.
rahul, kumarjit10r, anish@isid.ac.in